



Environmental Impact Assessment (EIA) Report for the Liquefied Natural Gas Project in Cabo Delgado

FINAL EIA REPORT- VOLUME I

Project ref: 0133576

February 2014





Document	Environmental Impact	Volume	FINAL EIA REPORT-
Title:	Assessment (EIA) Report	No:	VOLUME I & II
	for the Liquefied Natural		
	Gas Project in Cabo	Document	February 2014
	Delgado	Date:	
Project	0133576	Revision	September 2014
Ref:		Date:	

Sections / Chapters	Version	Change Description
Abbreviations/ Units	Portuguese	The unit "mg/l-1" was replaced by "mg/l or mg.l-1" to represent milligrams per litre.
Acronyms Chapter 9 Chapter 13	Portuguese	The Acronym "ASA" used to denote Environmental Health Area (Área de Saúde Ambiental) has been changed to EHA.
Acronyms Chapter 9 Chapter 13	Portuguese	The acronym "TS" used to denote "sex workers" (Trabalhadoras de Sexo) has been removed.
Acronyms	Portuguese	A Portuguese translation for the following acronyms has been included:
		 EGPAF FH GDACE HYCOM
Acronyms	Portuguese	Where an English acronym is used, the English definition has been provided in addition to the Portuguese definition.
Acronyms	Portuguese and English	The definition for the acronym "DWAF" now states: "South African Department of Water Affairs and Forestry".
Acronyms Chapter 2	Portuguese and English	The acronym "MdP" to denote "Ministry of Fisheries" has been replaced with "MPescas".
Acronyms	Portuguese	The word "manganésio" used to explain the chemical symbol "Mn" has been replaced with "manganês".
Acronyms	Portuguese	The term "non-aqueous based mud" used to explain WMB, has been replaced by "water based mud".
Acronyms	Portuguese	The acronyms LBA and LBS were deleted as the EIA Report uses WBM and SBM (for water based muds and synthetic based muds respectively) and not LBA and LBS.
Chapter 3	Portuguese	In <i>Table 3.4</i> , the date the EIA Report was submitted to MICOA has been corrected to state 2014, not 2013.

Environmental Impact Assessment (EIA) Report for the Liquefied Natural Gas Project in Cabo Delgado

FINAL EIA REPORT

February 2014

Project Ref: 0133576

Prepared by: ERM and Impacto

For and on behalf of ERM and Impacto
Approved by: Andrew Bradbury
Signed: Andrea franker
Position: Partner
Date: 24 February 2014

This report has been prepared by Environmental Resources Management the trading name of Environmental Resources Management Southern Africa (Pty) Limited, with all reasonable skill, care and diligence within the terms of the Contract with the client, incorporating our General Terms and Conditions of Business and taking account of the resources devoted to it by agreement with the client.

We disclaim any responsibility to the client and others in respect of any matters outside the scope of the above.

This report is confidential to the client and we accept no responsibility of whatsoever nature to third parties to whom this report, or any part thereof, is made known. Any such party relies on the report at their own risk.

This proposal is provided solely for the purposes set out in it and may not, in whole or in part, be used for any other purpose without ERM's prior written consent. This proposal may not, in whole or in part, be reproduced without ERM's prior written consent.



EIA Report - Contents

Volume I – Introduction, Project Description and Baseline

Non-Technical Summary (NTS) Abbreviations/Units, Acronyms and Glossary

- 1. Introduction.
- 2. Regulatory Framework.
- 3. The EIA Process, Approach and Methodology.
- 4. Project Description.
- 5. Consideration of Alternatives.
- 6. Baseline- Introduction and Geographical Context.
- 7. Environmental Baseline Offshore and Near Shore.
- 8. Environmental Baseline Onshore.
- 9. Socio-economic and Community Health Baseline.

Volume II – Impact Assessment, Management, Implementation and Conclusions

- 10. Introduction to Impact Assessment and Mitigation.
- 11. Offshore and Near Shore Environmental Impact Assessment and Mitigation.
- 12. Onshore Environmental Impact Assessment and Mitigation.
- 13. Socio-economic Impact Assessment and Mitigation.
- 14. Unplanned Events.
- 15. Cumulative Impacts.
- 16. Impact Assessment Summary.
- 17. Environmental and Social Management System and Environmental and Social Management Plan.
- 18. Conclusion.
- 19. References.

Volume III – Annexes

- A Public Participation Report.
- B MICOA Approval of EPDA and ToR.
- C Baseline Methodologies.
- D Tabulated ESMP.
- E Waste Management Plan.
- F Decommissioning and Rehabilitation Plan.
- G Baseline Support Material.
- H Emergency Response Plan.
- I Initial Resettlement Plan.



Volume I – Introduction, Project Description and Baseline

Non-Technical Summary (NTS) Abbreviations/Units, Acronyms and Glossary

- 1. Introduction.
- 2. Regulatory Framework.
- 3. The EIA Process, Approach and Methodology.
- 4. Project Description.
- 5. Consideration of Alternatives.
- 6. Baseline- Introduction and Geographical Context.
- 7. Environmental Baseline Offshore and Near Shore.
- 8. Environmental Baseline Onshore.
- 9. Socio-economic and Community Health Baseline.

NON TECHNICAL SUMMARY (NTS)

ABBREVIATIONS/ UNITS, ACRONYMNS AND GLOSSARY

1	INTRODUCTION	1-1
1.1	THIS DOCUMENT	1-1
1.2	THE PROJECT PROPONENT	1 - 1
1.3	Тне Ргојест	1-3
1.4	THE REQUIREMENT FOR AN EIA	1-7
1.5	ENVIRONMENTAL PRACTITIONER	1-7
1.6	STRUCTURE OF THIS REPORT	1-9
2	REGULATORY FRAMEWORK	2-1
2.1	INTRODUCTION	2-2
2.2	Mozambican Legal/Regulatory Framework	2-2
2.3	INTERNATIONAL CONVENTIONS	2-23
2.4	GOOD INTERNATIONAL INDUSTRY PRACTICE	2-26
2.5	REQUIREMENTS OF OTHER INTERNATIONAL INSTITUTIONS OF RELEVANCE	2-29
2.6	NUMERICAL STANDARDS AND GUIDELINES RELEVANT TO THE PROJECT	2-30
2.7	ANADARKO COMPANY POLICY	2-37
2.8	ENI'S COMPANY POLICY	2-38
3	THE EIA PROCESS APPROACH AND METHODOLOGY	3-3
3.1	OVERVIEW OF THE EIA PROCESS	3-3
3.2	OVERVIEW OF APPROACH	3-3
3.3	Phase 1: EIA Pre-Assessment Application (Screening)	3-5
3.4	PHASE 2: ENVIRONMENTAL PRE-FEASIBILITY STUDY AND SCOPE OF	
	DEFINITION (EPDA/ SCOPING)	3-6
3.5	Phase 3: Impact Assessment	3-9
3.6	EIA METHODOLOGY	3-1 5
3.7	MANAGEMENT AND MONITORING	3-21
3.8	DEALING WITH AND MANAGING GAPS AND UNCERTAINTY	3-22
4	PROJECT DESCRIPTION	4-1
4.1	INTRODUCTION	4- 1
4.2	OVERVIEW OF THE PROJECT	4-1
4.3	OFFSHORE PROJECT DESCRIPTION	4-9
4.4	ONSHORE PROJECT DESCRIPTION	4-2 4

4.5	NEAR SHORE PROJECT DESCRIPTION	4-42
4.6	MANAGEMENT OF EMISSIONS, DISCHARGES AND SOLID WASTES	4-53
4.7	DECOMMISSIONING AND CLOSURE	4-68
5	CONSIDERATION OF ALTERNATIVES	5-1
51	INTRODUCTION	5_1
5.2	Site Ai ternatives	5-1
53	I AVOLIT ALTERNATIVES	5-11
5.5 5.4	TECHNOLOCY AND PROCESS ALTERNATIVES	5-11
5.5	No-co Ai ternative	5-14
5.6	Consideration of Alternatives through FFFD	5-10 5-18
5.0		5-10
6	BASELINE - INTRODUCTION AND GEOGRAPHICAL CONTEXT	6-1
6.1	OVERVIEW AND STRUCTURE OF THE EIA BASELINE	6-1
6.2	SUMMARY	6-1
6.3	Sources of Information	6-3
6.4	GEOGRAPHICAL SETTING	6-4
6.5	Areas of Influence and Study Area	6-7
6.6	CLIMATE AND METEOROLOGY	6-10
6.7	AIR QUALITY	6-17
6.8	Noise	6-21
6.9	LANDSCAPE AND SEASCAPE CHARACTER AND VISUAL AMENITY	6-24
6.10	NATURE CONSERVATION	6-42
7	ENVIRONMENTAL BASELINE - OFFSHORE AND NEAR SHORE	7-1
7.1	INTRODUCTION	7-1
7.2	Summary	7-1
7.3	Sources of Information	7-3
7.4	Regional Context – Quirimbas Archipelago	7-3
7.5	OFFSHORE ENVIRONMENT – PHYSICAL CONDITIONS	7-6
7.6	OFFSHORE ENVIRONMENT – MAJOR BIOLOGICAL FEATURES	7-22
7.7	NEAR SHORE ENVIRONMENT – PHYSICAL CONDITIONS	7-37
7.8	NEAR SHORE ENVIRONMENT – MARINE HABITATS	7-59
7.9	NEAR SHORE ENVIRONMENT – MAJOR BIOLOGICAL FEATURES	7-95
8	ENVIRONMENTAL BASELINE – ONSHORE	8-1
8.1	INTRODUCTION	8-1
8.2	Geology and Terrain	8-3
8.3	Soils and Land Capability	8-5
8.4	GROUNDWATER	8-19
8.5	Hydrology	8-33
8.6	SURFACE WATER ECOLOGY	8-44
8.7	VEGETATION	8-112

8.8	Herpetofauna	8-132
8.9	AVIFAUNA	8-151
8.10	MAMMALS	8-180
8.11	SUMMARY OF KEY ONSHORE ENVIRONMENTAL SENSITIVITIES	8-200
9	SOCIO-ECONOMIC AND COMMUNITY HEALTH BASELINE	9-1
9.1	INTRODUCTION	9-1
9.2	SUMMARY	9-1
9.3	Sources of Information	9-7
9.4	SOCIO-ECONOMIC STUDY AREA	9-8
9.5	CABO DELGADO PROVINCE: ADMINISTRATIVE AND POLITICAL OVERVIEW	9-9
9.6	CABO DELGADO PROVINCE: SOCIO-DEMOGRAPHIC INDICATORS	9-10
9.7	CABO DELGADO PROVINCE: ECONOMIC ACTIVITIES	9-16
9.8	PALMA DISTRICT POLITICAL AND ADMINISTRATIVE ORGANISATION	9-35
9.9	PALMA DISTRICT AND AFUNGI PROJECT SITE: SOCIO-DEMOGRAPHIC	
	INDICATORS OF THE AREA OF DIRECT INFLUENCE	9-37
9.10	PALMA DISTRICT AND AFUNGI PROJECT SITE AND SURROUNDS: USE OF	
	NATURAL RESOURCES	9-50
9.11	PALMA DISTRICT AND AFUNGI PROJECT SITE AND SURROUNDS: LAND USE	
	AND OCCUPATION PATTERNS	9-52
9.12	PALMA DISTRICT: NON-GOVERNMENTAL ORGANISATIONS AND DONORS	9-59
9.13	PALMA DISTRICT AND AFUNGI PROJECT SITE SOCIAL SERVICES	9-61
9.14	PALMA DISTRICT AND AFUNGI PROJECT SITE ECONOMIC ACTIVITIES AND	
9.15	AFUNGI PROJECT SITE AND SURROUNDS (INCLUDING SENGA AND	
	MAGANJA) CULTURAL AND RELIGIOUS PROFILE	9-131
9.16	AFUNGI PROJECT SITE AND SURROUNDS (INCLUDING SENGA AND	
	MAGANJA): EXPECTATIONS REGARDING THE PROJECT	9-134
9.17	Shipping and Navigation	9-139
9.18	Archaeology and Cultural Heritage	9-14 8



Non-Technical Summary

Mozambique Gas Development Project

Non-Technical Summary (NTS)



Project Ref: 0133576 February 2014







Mozambique Gas Development Project

Introduction and Background

Natural gas discoveries made in the Rovuma Basin off the coast of northern Mozambique are among the world's most significant discoveries in the last 20 years. Anadarko Moçambique Área 1, Lda (AMA1) holds rights to explore, develop and produce natural gas reserves in Area 1 in the Rovuma Basin and Eni East Africa S.p.A (eni) holds similar rights to explore, develop and produce in Area 4. These areas are adjacent to one another and a number of gas reservoirs or gas fields have been discovered within each area. AMA1 and eni are joint proponents for the Project, working to develop a Liquefied Natural Gas (LNG) Facility and associated infrastructure to convert the natural gas to a liquid state and export it to international markets.

The proposed LNG Project (hereinafter "the Project") would see Mozambique become one of the world's leading natural gas exporting countries. Figure 1 below illustrates the location of the Project.

The Project is required to obtain permission (in the form of an environmental licence(s) from the Government of Mozambique) before AMA1 and eni can construct and operate the facilities. An Environmental Impact Assessment (EIA) Report has therefore been developed to inform the Government's decision making process. The EIA Report documents the process undertaken for identifying, assessing, mitigating and managing the biological, physical and socio -economic impacts of the Project.



Figure 1: Location Area

The Project Developers

AMA1

AMA1 is a wholly-owned subsidiary of Anadarko Petroleum Corporation (APC) and is a commercial entity duly registered under the Republic of Mozambique Laws. AMA1 has offices in Maputo and Pemba. APC is headquartered in The Woodlands, Texas, United States of America and is among the largest independent oil and natural gas exploration and production companies in the world. Information on AMA1 can be accessed on the internet at:

http://www.anadarko.com/Operations/Pages/ LNGmozambique.aspx

ENI

eni S.p.A is an Italian company with activities in close to 85 countries. Eni East Africa S.pA (eni) is an affiliate of eni S.p.A. and has a branch in Mozambique with offices in Maputo and Pemba. eni S.p.A. is one of the leading global operators in the deep-water sector worldwide and is currently involved, as operator or partner, in exploration activities in deep water in the Gulf of Mexico and along the coast of Brazil. More information on eni can be accessed on the internet at:

http://www.eni.com/en_IT/media/casebook/casebook/ mozambigue.html



Economic Benefits of the Project



The initial LNG development (two LNG trains estimated 10mtpa) represents an overall investment of up to US\$25—30 billion, making the project the largest single investment project in Mozambique to date. Assuming export of the LNG to premium markets, such as Japan and the Far East, the Government of Mozambique should see a significant increase in Gross Domestic Product (GDP) and increased revenue through its royalty, tax and equity gas rights.

Mozambique will benefit from a substantial increase in Government revenue for several decades. This economic benefit could be used to improve the health, education and quality of life of the people of Mozambique.

As the number of trains and volume of LNG export, increases, the economic benefits could be compounded several times. The Government of Mozambique has stated it wishes to encourage industrial development using natural gas in the area of the Project. This could further increase the social and economic benefits derived from the Project.



Project Description

The purpose of the Project is to gather, process, and export natural gas in liquid form known as LNG. This LNG will be used as a fuel source in other countries.

The process begins offshore in Area 1 and Area 4 of the Rovuma Basin, where natural gas will be extracted via subsea wells from gas reservoirs (storage areas) up to 1,500m below the seafloor. The collected gas will be transported to the onshore LNG facility by pipelines on the seafloor. Once onshore, the gas will be processed in the LNG facility to remove impurities, converted to liquid (by cooling the gas) and stored in specially designed storage tanks.

The LNG will then be transported through pipelines to an export jetty where it will be loaded into specialized LNG carriers to be transported to international markets. These specially designed ships maintain the LNG in a refrigerated liquid state for sea voyages of several thousand kilometres. The Project has an initial 30 year lifespan but this may be extended depending on future gas reserve development.

The three main elements of the Project are as follows and illustrated in Figure 2:

- Offshore drilling of wells and installation of pipelines on the seafloor to connect the wells and then bring the natural gas to the shore.
- Near Shore construction of a Multi-Purpose Dock and jetties. The dock will house support vessels and allow for equipment and material (for onshore construction) to be brought ashore. LNG carriers will berth at the jetties while they are filled with the LNG.
- **Onshore** construction and operation of the LNG Facility and all associated infrastructure eg housing, construction camp, airport etc.

The above three main elements are described in detail overleaf.

The project description will be refined through the Front End Engineering and Design (FEED) process currently being undertaken). Should there be any significant departure from the project description in the EIA, such that impacts are different or there needs to be new mitigation measures, the Project will undertake the necessary environmental processes (eg addendum to the EIA).

Figure 2: Illustrative Scheme of Project





Dffshore

The term 'Offshore' refers to the deep water environment. Offshore Project components comprise the wells, infrastructure connecting the wells with systems known as manifolds, to the gas pipelines leading up to Palma Bay (areas within Palma Bay are referred to as the Near Shore). The process begins in Area 1 and Area 4, where natural gas will be extracted via subsea wells from gas reservoirs below the seafloor. The Offshore Project components will consist of production wells and the pipeline system. Production wells are planned for AMA1's Golfinho and Prosperidade gas fields and for eni's Mamba gas field. Up to 60 production wells are planned for the Prosperidade and Mamba gas field and a further 60 production wells are planned for the Golfinho gas field.

The wells in each field will be connected to central collection systems on the sea bed so gas from multiple wells can be collected together. The collected gas will be transported by pipelines (running in a single corridor as they approach the near shore) from the offshore facility to the onshore facility. The offshore system is anticipated to produce up to 4 billion cubic feet of gas a day.

The offshore facility will be constructed by the drilling and installation of wells using dynamically positioned drill ships. Multi-purpose vessels will also be used to install the offshore central collection systems on the seafloor. Once operating, gas will be collected and transported to the onshore facility.

Figure 3 : Example of a Drill Ships and Construction Vessels



Example of a Deepwater Pipelay Vessel



Example of a Flexible Pipe and Umbilical Lay Vessel

Example of a Light Construction Vessel











The Near Shore refers to the shallow water within Palma Bay; from the islands of Tecomaji and Rongui to the shoreline. The Near Shore Project components will consist of port facilities (jetties and berths), a Multi-Purpose Dock to support construction activities (eg bringing in heavy equipment and materials), a shipping lane and the pipeline corridor within Palma Bay. Figure 4 below provides an indicative illustration of the near shore infrastructure.

Access channels to the port facilities will be widened and deepened by dredging the seabed in Palma Bay to allow large ships to reach the jetty and berths. Dredging will also be required in the Near Shore to enable pipe-laying. Some dredged material will be used for infill in the near shore while the rest will be disposed of at designated areas offshore.









Onshore

The precise layout of the various Onshore components will be defined during the current Front End Engineering and Design (FEED) process. Thus indicative layouts have been used as the base case of the impact assessment. Figure 5 provides an indicative illustration of the Onshore Project components. Vegetation will need to be cleared and the land leveled prior to construction of the Onshore Project footprint. The Project footprint area is currently approximately 3,600 ha, within the allocated approximately 7,000 ha DUAT area.

The Onshore Project components comprises facilities necessary to treat and process natural gas and supporting infrastructure such as worker accommodation facilities, construction areas, access roads and an airport. The LNG processing plant will receive gas from the offshore facilities, remove impurities in the gas and produce LNG by progressively cooling the gas to a temperature of -163°C. The LNG will be stored in cooled storage tanks on site and transferred by pipeline to the jetties for loading onto ships. The processing plant will produce between 3,000 and 5,000 barrels of LNG a day. The process will also generate water that will be recycled into the process and natural gas condensate, a byproduct that will be sold.

Figure 5: Illustration of Onshore Infrastructure



6





The EIA process

The purpose of the EIA is to predict the significance of the Project's impacts on the existing (baseline) physical, biological and socio-economic environment; and to identify measures to minimise negative impacts and maximise positive ones. This information will be used to inform decision-making by the Government of Mozambique.

The EIA study comprised the following steps:

- An integrated environmental, socioeconomic and engineering Site Selection study was undertaken to determine the most appropriate site for the construction and operation of the LNG Facility;
- The potential impacts of the Project were initially identified during the 'Estudo de Pré-Viabilidade Ambiental e Definição de Âmbito' (EPDA) or Scoping Phase;
- The existing baseline conditions and any environmental or socio-economic sensitivities were determined through fieldwork and review of existing information;
- Stakeholder concerns raised during the EPDA Phase and Impact Assessment Phase were considered;
- The significance of impacts prior to mitigation measures being applied was assessed;

 The significance of the impacts after and the model of the project are determined (by project)
 The baseline conditions are determined (by project)

 Windows and the impacts after a set of the impacts after a set o

 Mitigation measures were developed and refined during workshops and meetings with the Project engineers and contractors:

Specialist experience and knowledge coupled with modelling in some cases (eg noise, drilled cuttings dispersion) was used to inform the impact assessments;

- Workshops and meetings were convened to ensure that the proposed mitigation measures were practical and implementable;
- The Onshore Project layout was revised and reduced based on identified baseline sensitivities, outcomes of the impact assessment and following inputs from environmental and social specialists;
- The significance of residual impacts, prior to mitigation measures being applied was assessed; and

Figure 6: EIA Process Overview

• An Environmental and Social Management Plan (ESMP), Waste Management Plan and Decommissioning and Rehabilitation Plan were developed and a process for training and educating staff about these plans was prepared.

Stakeholder Engagement

Stakeholder engagement (including communication with local communities) is a critical part of the EIA process and activities were carried out at key stages of the EIA process to ensure that stakeholder concerns and comments are addressed in the EIA. Meetings were held with stakeholders during the EPDA and Impact Assessment Phases. In addition, focus group meetings were held during the Impact Assessment Phase with key stakeholders. The Draft EIA Report was made available for public comment between 27 August and 31 October 2013.

All comments received throughout the engagement process and commenting period were recorded, and responses are provided in a Public Participation Report (*Annex A* of EIA Report).

The Draft EIA Report was updated based on comments received prior to submission to MICOA in February 2014.





Impacts were identified by the EIA team with inputs by stakeholders (eg local communities, government departments and local communities). An impact is any change to a resource or receptor brought about by the presence of the Project component or by the execution of a Project-related activity. The impacts assessed fall into two main categories; environmental and socio-economic. Environmental impacts include both physical impacts of the Project such as changes in air quality and biological impacts such as changes to marine and land based habitats.

The socio-economic impacts are effects of the Project on people and their livelihoods. This includes positive impacts such as employment or increased demand for local goods and services as well as negative impacts such as loss of access to farm land or access to parts of Palma Bay for engaging in subsistence livelihoods activities. The following aspects were considered when determining the significance of identified impacts:

- nature of impact: positive or negative;
- duration of impact: temporary, short term, medium term, long term or permanent;
- scale of impact: onsite, local, regional, national or international/ transboundary; and
- intensity of the impact: negligible, low, medium, high.

Table 1 shows the definitions for the categories of significance used in this EIA.

Table 1 : Definition of Significance Levels		
NEGLIGIBLE	The existing environmental and social conditions will not be affected or the affect is not detectable. A negligible impact is likely to be of no concern to the government, communities and organisations.	
MINOR	The environmental and/or social conditions will be affected, but the impact small enough that it is unlikely to be of concern to the government, communities and organisations.	
MODERATE	An impact of moderate significance is one within accepted limits and standards. The emphasis for moderate impacts is on demonstrating that the impact has been reduced to a level that is as low as reasonably practicable (ALARP).	
MAJOR	A major impact is one where there will be a large change to communities or the natural environment. The aim of the EIA is to provide ways to stop these large impacts from occurring. At times this is not possible and it is up to the government to decide if this is acceptable when considering the benefits of the Project.	

Mitigating Impacts:

The Project philosophy regarding mitigation is outlined in Table 2. The Project first sought to avoid impacts to the extent practical. If impacts could not be avoided they were mitigated through modifying the design or implementing mitigation at the impact site. In cases where these first two mitigation strategies were not possible, the Project would aim to remediate impacts wherever possible and finally investigate options for compensation or offsets where necessary or required.

Table 2 : The Mitigation Hierarchy for Planned Project Activities
AVOID AT SOURCE; REDUCE AT SOURCE Avoiding or reducing at source is 'designing' the project (embedded controls) so that a feature causing an impact is designed out (eg a waste stream is eliminated) or altered (eg reduced waste volume). Often called minimization.
ABATE ON SITE This involves adding something to the basic design to abate the impact - pollution controls fall within this category. Often called 'end-of-pipe'.
ABATE AT RECEPTOR If an impact cannot be abated on-site then measures can be implemented off-site. An example of this would be to use a silt trap to capture any sedi- ment that may flow into streams.
REPAIR OR REMEDY Some impacts involve unavoidable damage to a resource, eg land disturbance. Repair involves restoration and reinstatement measures, such as re- vegetation
COMPENSATE IN KIND Where other mitigation approaches are not possible or fully effective, then compensation, in some measure, for loss, damage, and general intrusion might be appropriate.



Key Impacts for Decision Making

The EIA has assessed the potential impacts of the proposed Project on the biophysical and socio-economic environment. Throughout the EIA process evolving project information has been fed into the assessment process, allowing for the EIA to be updated as the Project concept was developed. Thus, impacts have been assessed taking into account the mitigation measures that were built into project design. Potential 'cumulative' impacts (ie impacts acting in conjunction with each other on a common resource or receptor) and the risk of accidents (eg gas pipeline rupture) have also been assessed as an integral part of the assessment exercise.

The findings of the EIA process are presented in this section. A large number of potential impacts were assessed in the EIA process and the detailed findings are presented in the EIA Report.

BASELINE

The offshore project area is located in deep waters, in depths of around 1,000m to 2,300m. The offshore marine environment supports a wide range of larger animals such as whales and dolphins as well as many species of fish, turtles and sea birds. Deep water reef structures (mainly hard substrate structures) have been observed on the seabed.

The near shore environment, in and around Palma Bay, has clear water (low levels of suspended sediment), temperatures in the range of 30 to 35°C and a number of different habitats; rocky bottom, sand beaches, mangroves and mudflats. Figure 7 shows a range of the typical habitat types. Palma Bay supports sea grass beds close to the shore with coral reefs scattered within the bay and around the islands of Tecomaji and Rongui.

PROJECT ACTIVITIES

Project activities that could affect this baseline include dredging, disposal of dredge material, installation of subsea infrastructure, and the construction of the jetties and Multi-Purpose Dock.

Figure 7 Habitats in the Offshore and Near Shore Environment





Impacts due to discharge of drill cuttings:

Dispersion modelling of drill cuttings and the sensitivity of offshore benthic fauna and flora indicates that impacts from physical inundation (burial and change in sediment grain size) will be of MINOR significance before and after mitigation. The drilling of the offshore production wells will be the primary activity likely to result in potential impacts to water quality and marine ecology (such as whales, dolphins and benthos) in the offshore area. With the implementation of mitigation measures, all residual impacts are predicted to be of NEGLIGIBLE and MINOR significance in the offshore marine environment.

Deepwater reef structures and associated organisms, however, may be at risk of enduring more severe effects. This is because the recovery of reef structures is expected to take an appreciable amount of time due to generally slow growth rates of these organisms. Therefore the predicted impact to offshore reef communities may potentially be of MODERATE significance prior to mitigation. It should be noted that both the high and low-relief structures observed in the deep waters offshore largely comprise scattered rock and sediment, and do not apparently support high densities of fauna. With the implementation of mitigation, this impact will be reduced to MINOR.

Impacts due to discharge of residual muds:

Impacts on benthic marine organisms or those in the water column (ie plankton) from discharges of residual muds (small amounts of muds that remain on the drill cuttings after treatment) are predicted to be of MINOR significance because of low toxicities of the quantities and types of muds used/or discharged to the offshore environment. Impacts will be reduced to NEGLIGIBLE with the implementation of mitigation.

Impacts due to discharge of hydrotest water:

The impacts on marine ecology and/or marine ecological processes associated with the discharge of hydrotest water at depths of approximately 1,500m in the offshore gas fields during commissioning (ie the construction phase) will be NEGLIGIBLE, as the hydrotest water will be discharged in a phased approach, at pressures that will ensure that the water quality effects of the discharges are restricted to close proximity of the release points.

Impacts due to increased traffic:

Potential impacts from vessel and helicopter noise, lighting and movements on offshore marine ecology (birds, fish, benthic fauna etc) with the exception of marine mammals will be NEGLIGIBLE. Effects of vessel collisions or disturbance on whales may be more severe due to their conservational value or importance and the associated impact is predicted to be of MODERATE significance. However, mitigation measures specific to whales will ensure that vessel collisions with whales are avoided and thus impact significance is reduced to NEGLIGIBLE.

Impacts due to habitat modification:

The introduction of the subsea infrastructure, a hard structure, on to the seabed in the Offshore Project Area will result in changes to the character of the sea bed and consequently the diversity and structure of the benthic community. Such an impact will be of MODERATE significance, particularly at deep water reef structures, where benthic organisms including deepwater reef structures, may be affected. Mitigation measures will ensure that the subsea infrastructure is located to avoid areas of sensitive benthic communities to the extent practical. Thus reducing the impact significance to NEGLIGIBLE.



Summary of Key Near Shore Environmental Impacts

Construction activities in the Near Shore are likely to result in disruptions to seagrass, coral reefs and some loss of mangrove and estuary areas. These features play critical roles in ecological interrelationships and directly and indirectly support productivity and biodiversity in the Palma Bay area. The implementation of mitigation has helped to reduce the significance of such impacts but the potential remains for bay-scale productivity to be affected.

Impacts due to dredging: Pre-mitigation impacts to marine ecology from dredging activities in the Near Shore Project Area have been assessed as severe, with a number of MAJOR significant impacts being identified. Examples of such impacts include; the effects on seagrass, coral and associated biological communities from increased turbidity in the water column, cutting a trench through coral reef and rock, depositing fine sediment on benthos and modifications to the seabed. The significance of these MAJOR impacts is largely reduced through mitigation. Key mitigation measures include changing the proposed dredging/cutting techniques and avoidance of areas considered particularly sensitive (ie corals along the pipeline route) and measures to reduce turbidity.

The deposition of fine sediments from dredging activities can inundate seagrass beds and also cover corals with a veneer of sediment that may take considerable time to recover. The re-suspension of sediment and subsequent deposition is unavoidable and subsequent effects to benthos and coral are likely to remain as MAJOR significance post-mitigation.

Dredge material discharged in a dredge placement area at the head of the Afungi Canyon in Palma Bay will smother and possibly result in negative impacts to benthos within the designated 1km² placement area. The impact will be of MODERATE significance within dredge placement area itself pre-mitigation, and MINOR post-mitigation. Benthos will likely recover within 1 to 3 years after the cessation of dredging.

Impacts due to modification of the beach: The installation of Near Shore infrastructure across the intertidal beaches and extending into the shallow subtidal zone will modify beach structure and dependant ecological processes. This is likely to result in a loss of parts of the productive sand beach and subtidal zones and areas of seagrass meadow. It will allow the establishment of hard substrate communities in the lower intertidal and corals, sponges and associated organisms in the subtidal and may also facilitate colonisation by alien and potentially invasive species. The impacts to the marine biotopes and associated communities within the bay will be MODERATE for the construction and operational phases. With mitigation, including design mitigation, these impacts will be reduced to MINOR significance.

Impacts due to increased noise: Impacts to fish, whales, dolphins and turtles from noise associated with pile driving in Palma Bay are predicted to be of MODERATE significance although the extent of the effects differs among groups. Should a 'soft start' procedure be implemented prior to construction activities when megafauna are present in the bay, the impact will be reduced to MINOR.

Impacts due to alien invasive species: If invasive alien species are transferred into Palma Bay via ballast water, effects on biodiversity and marine ecology, including seagrass and corals may be of MODERATE significance. The progressive adoption of ballast water control measures and processing techniques as sanctioned by International Maritime Organisation (IMO) will further reduce the likelihood of releasing of non-indigenous organisms. However, due to the high magnitude of the effects that may arise should invasive species become established in northern Mozambique, the significance rating would remain as MODERATE.

Impacts due to discharges into the bay: Discharges from the proposed desalination and sewage treatment plants as well as from treated produced water and stormwater from the LNG Facility is are likely to have NEGLIGIBLE to MINOR significant impacts on near shore water quality and marine flora and fauna in Palma Bay prior to mitigation. All impacts from sources of discharge post-mitigation will be NEGLIGIBLE.

Impacts due to waste discharges: Discharged solid and liquid wastes from marine vessels during any phase of the Project could potentially result in impacts of MODERATE significance through the proliferation of litter and compromised water quality harming marine organisms, seabirds and biodiversity in Palma Bay. With effective mitigation of waste with appropriate facilities impacts will be reduced to NEGLIGIBLE.

Impacts due to loss of an estuary and its mangroves: The impacts on marine ecology resulting from the loss of the estuary and associated mangrove stand to the east of Afungi Project Site during the construction phase will be of MODERATE significance. The impact will remain as MODERATE significance post-mitigation given the loss of the multi -species mangrove stand.

Impacts due to the security exclusion zones: The establishment of security (exclusion) zones around the LNG Facility and the Near Shore Project infrastructure during construction and operations will displace artisanal fishing effort and locally increase exploitation pressure on the fish community outside of the zones resulting in impacts of MINOR significance. Mitigation measures to help reduce such impacts have yet to be identified.



Summary of Key Onshore Environmental Impacts

BASELINE

The onshore project area is located in an area that has three main habitat types, marshlands, wetlands and woodlands. All of these habitats are considered important due to the animal and plant life they support. Figure 9 shows some of the plants and animals in the Afungi Project Site. The wetlands are fed by drainage channels upstream of the site; the flows in these channels are highly variable with very high flow in the wet season and very low flow in the dry season. The wetlands are of particular importance as they provide habitat for frogs and lizards (amphibians). These amphibians are eaten by other animals and largely support the food chain in the area.

Figure 8 Examples of plants and animals in the Afungi Project Site





PROJECT ACTIVITIES

During the construction phase, the activities that will impact on the baseline are the clearance of vegetation, infilling of an estuary, and other general site preparation activities (eg levelling of land). During the operation phase, activities that could affect the baseline are associated with accidental spills, runoff and sedimentation.

REVISION TO ONSHORE PROJECT LAYOUT

Field studies conducted during the Baseline Phase of the EIA identified and mapped sensitive habitats for a variety of vegetation, terrestrial fauna and avian species. Once these species-specific sensitive habitat maps were overlaid (see Figure 9), it became apparent that certain areas within the Afungi Project Site were more sensitive than others. Most of the species relied heavily on the wetlands within the Onshore Project Footprint Area for the ecological functions they provide (food, water, breeding habitat, etc).

Through baseline studies and sensitivity mapping, it became obvious the Project could reduce adverse environmental impacts by revising the Onshore Project Footprint Area. With the sensitivity map in mind, the Project began to investigate methods to avoid or minimise potential footprint impacts. This was facilitated by a series of interactions held between the EIA and Project Engineering Teams. The EIA Team, in conjunction with AMA1, revised the base case Project layout (Figure 9) to avoid or minimise impacts on the identified high-sensitivity areas. Figure 10 shows the Revised Project Footprint Area. The potential FEED Contractors were then tasked with determining whether they could design the Project to work within the revised areas. Each of the potential FEED Contractors confirmed that they were able to work within the revised layout.

This mitigation exercise enabled the avoidance of some impacts and the minimisation of others to as low as reasonably practicable (ALARP). The Revised Project Footprint Area effectively reduces the disturbance of areas classified as High to Very High terrestrial ecological sensitivity from approximately 2,340ha to 1,695ha, thereby avoiding the disturbance of approximately 645ha of High to Very High sensitivity.





Project Footprint Area - Base Case and Revised

Revised Footprint Area

The Onshore Project Footprint (see Figure 9) was revised to minimise impacts to area considered to be of high sensitivity to habitats and various animals such as birds, mammals and frogs. This was a key mitigation measure developed and the revised Project footprint is shown below in Figure 10.





Figure 10 Revised Footprint







Villages / Settlements

Regional Roads

Legend





Summary of Key Onshore Environmental Impacts

Impacts due to air emissions: Air dispersion modelling undertaken indicates that during the operational phase of the Two-Train LNG Facility, there will be minimal impacts to human and ecological receptors outside the Afungi Project Site from changes to air quality. Impacts to receptors outside of the Afungi Project Site will be of NEGLIGIBLE significance for all pollutants modelled pre- and post- mitigation with the exception of SO₂ which will be of MINOR significance prior to mitigation and NEGIGIBLE post-mitigation.

However, with the expansion of the LNG Facility to six trains, a potential impact of MODERATE significance may occur from increased levels of SO₂ pre-mitigation. Further consideration of the design of the LNG Facility (in terms of both stack height and turbine technology adopted) is recommended to avoid the potential of exceeding air quality standards during operations of an expanded six train facility.

Impacts due to emissions of greenhouse gasses: Assuming Mozambique's Greenhouse Gas (GHG) emissions will increase by 8 percent per year (based on World Bank GDP predicted growth figures), emissions from the Project will increase the country's global contribution of GHG emissions from between 0.4 percent per year up to 10 percent per year, depending on the year and period of development (ie construction or operation). The pre-mitigation impact of the Project on Mozambique's national GHG emissions is likely be of MAJOR significance both during the construction and operational phases of the Project. Given the scale and nature of the Project, while good practice can be employed to reduce the GHG emissions, the overall significance of the impact is not expected to significantly change post-mitigation.

Impacts due to noise generation: Noise modelling has shown that during normal construction and operational activities noise levels will remain in compliance with Mozambican and IFC noise limits/standards at Noise Sensitive Receptors at night and during the day. Impacts are therefore predicted to be of NEGLIGIBLE to MINOR significance premitigation and similar post mitigation.

Impacts to the visual landscape: Project activities during the construction phase, including the movement of vessels in and out of Palma Bay and the presence of highly visible equipment and machinery onshore, are likely to have a visual impact of MODERATE to MAJOR significance on Palma Bay Seascape Unit (ie in the coastal area of Palma Bay). During the operational phase this impact will increase to MAJOR significance, with the introduction and long term presence of the LNG Facility and associated infrastructure on Afungi Peninsula and within Palma Bay. Further south in the Afungi Peninsula to Cabo Nondo Seascape Unit, visual impacts experienced as a result of the Project will be of MINOR significance during construction and MODERATE during operations.

Visual impacts at a number of viewpoint locations in the wider area which are considered to be particularly sensitive to change are mainly expected to be of MINOR to MODERATE significance during construction. During operations when Project infrastructure is more visible, significance of visual impacts is likely increase to MODERATE to MAJOR, especially in nearby areas such as Palma town, Maganja and within Palma Bay. However, areas further removed from the Afungi Peninsula (Quiwia and the islands of Tecomaji and Rongui) visual impacts are expected to be of MINOR to MODERATE significance during operations. Vamizi Island and Olumbe are expected to have MINOR to NEGLIGIBLE visual impact during operation due to their distance from the Project.

Impacts on soil: Impacts on soil and land capability will predominantly be felt during the construction phase during site clearance activities, and will include soil compaction and topsoil loss, water and wind soil erosion and alteration of natural drainage. Impacts of MODERATE significance will be reduced to MINOR with the implementation of appropriate mitigation measures. Spills or leaks of fuel are likely to have NEGLIGIBLE impacts on soil resources as procedures will be put in place to prevent, contain, clean-up and dispose of any spillage.

Impacts due to groundwater use: Potential impacts to groundwater will be predominantly be felt during the early stage of the construction phase when groundwater will be the only source of fresh water. Groundwater abstraction from the production water supply wells may result in the lowering of groundwater levels (drawdown) in and around abstraction boreholes; however this is deemed to be of low magnitude potentially having a NEGLIGIBLE impact to community wells in the area adjacent to the production water supply wells. Additionally, water abstraction for Project use may have a NEGLIGIBLE impact on environmental receptors reliant on surface water.

Impacts on surface water ecology: The loss of wetland and estuarine habitat during the construction phase has been assessed as being of MAJOR significance. This significance rating is due to the loss of the ecological functions provided by the wetlands and of important habitat for species of conservational value or importance. The reconfiguration of the Project layout will result in a much reduced area of wetland and estuarine habitat being lost and this coupled with the additional mitigation measures proposed will reduce impact significance to MODERATE.

With the implementation of mitigation, all other impacts to wetland and estuarine flora and fauna are predicted to be MINOR, reduced from MODERATE in most instances, with the exception of impacts associated with increased turbidity and changes to sediment patterns. Increased turbidity in the wetland and estuaries is mainly likely to occur during the construction phase from onshore activities (eg removal of vegetation, site levelling, infilling of wetlands) and activities in the Near Shore (eg dredging). A change in sedimentation patterns may affect connectivity of the wetlands, primary production, benthic invertebrates, fish species etc. Impact significance could potentially be MAJOR during this phase of the Project but reduced to MINOR to MODERATE with mitigation.



Summary of Key Onshore Environmental Impacts

Impacts on vegetation: Much of the existing vegetation within the Afungi Project Site has been altered from its natural state (eg for subsistence agriculture) and fragmented. Site clearance activities during the construction phase are expected to result in further fragmentation and the removal or disturbance of some sensitive vegetation units. Site clearance is expected to result in an impact of MODERATE significance. Given the mitigated Revised Project Footprint layout, the areas of sensitive vegetation units to be lost will be reduced and impact significance is thus reduced to MINOR.

Impacts of MODERATE significance to ecological system functioning may occur due to the introduction of undesirable plant species during the construction phase, although with appropriate mitigation such impact will be reduced to NEGLIGIBLE.

Impacts on reptiles and amphibians: The impact of site clearance activities during the construction phase, in particular the infilling of wetland areas within the Afungi Project Site will be of MODERATE to MAJOR significance to reptiles and amphibians given the importance of freshwater wetlands in the functionality of their communities. However, the Revised Project Footprint Area reduces the amount of wetland disturbance and reduces this impact to MINOR significance.

Other impacts to reptiles and amphibians likely to arise during the construction and operational phases of the Project include mortality, disturbance and displacement due to the development of access roads and similar linear structures. Changes in water quality of the wetlands within the Afungi Project Site and adjacent areas also present potential impacts of MINOR to MODERATE significance to reptiles and amphibians. These impacts are reduced to NEGLIGIBLE to MINOR post-mitigation.

The influx of people into the wider area as a result of the Project is expected to put pressure on reptiles and amphibians population density through increased bush fires, reduced water quality from poor sanitation, subsistence poaching and hunting. Such activities could potentially result in reduced breeding success and depleted local populations of reptiles and amphibians and equating to impacts of MAJOR and MODERATE significance during the construction and operational phases respectively. Impacts to areas outside the Project's control are difficult to manage. However with the implementation of proper mitigation, the impact significance will be reduced to MODERATE and MINOR to MODERATE significance respectively.

Impacts on birds: Areas of important avian sensitivity are largely associated with estuarine salt marshes, freshwater wetlands, large intact forests and the inter-tidal zone and mangrove forests. Five International Union for Conservation of Nature (IUCN) listed bird species are known to occur within the Afungi Project Site and the loss of such bird habitat is considered MODERATE to MAJOR in significance. However the revision of the Project layout avoids or minimises impact to these sensitive areas. This in conjunction with additional mitigation measures serves to reduce the significance of the impact to MINOR to MODERATE.

Population influx may include the introduction of domestic animals (livestock and pets) into the area and the introduction of feral animals (dogs and cats). Feral animals are likely to prey on birds and livestock is likely to reduce habitat quality. As a consequence, avian impacts of MODERATE to MAJOR significance are possible during construction (when workforce is greatest); the impact significance is anticipated to be reducing to MODERATE during operations. With the introduction of mitigation measures, the impact will be reduced to MODERATE and MINOR to MODERATE during the construction and operational phases respectively. Other impacts to birds are reduced to NEGLIGIBLE to MINOR postmitigation.

Impacts on mammals: Project activities likely to result in impacts of MAJOR significant to mammals are largely associated with site clearance activities. Burrowing mammals, tree dwelling species and smaller mammals are likely to be most affected as they are less able to escape construction equipment or vehicles than larger species. Impacts during the operational phase are likely to be MINOR to MODERATE. However with management through mitigation, impacts are reduced to MINOR significance during both development phases.

The presence of infrastructure including roads, fencing, trenches and pipelines will cause loss or fragmentation of mammalian habitat. These may act as deterrents to mammals and have the potential to affect mammalian movements and cause disruption of mammalian migration. Impacts are assessed to be of MODERATE significance during both the construction phase and operational phase of the Project but reduce to MINOR with mitigation.

The most significant potential impacts to mammals remaining following mitigation are the indirect impacts associated with the influx of people into the area. These impacts are likely to be similar to those described for other species above. The various mitigation measures to be implemented by the Project are likely to reduce mammalian impacts to MODERATE during construction and MINOR to MODERATE during operations. All other impacts to mammals will be reduced to MINOR post-mitigation.



BASELINE

The Afungi Project Site is located within Palma District. Most people over the age of 15 in the district have received no formal education and illiteracy is high in the area. Formal employment is scarce and almost non-existent within the private sector. The majority of the population in the Afungi Project Site are therefore highly dependent on natural resources such as fishing and small scale agricultural activities. Figure 11 shows a local community's dependence on natural resources.

There are very few health care facilities in Palma District yet communities are regularly at risk of health problems arising from water and sanitation. The communities largely use natural water sources such as open wells and streams. The sanitation in the area is poor with few formal toilets, this puts the water sources at risk with potential for out breaks of diarrhoea and cholera.

PROJECT ACTIVITIES

The main Project activities that could change socio-economic conditions include the removal of access to land on the Afungi Peninsula, removal of access to parts of Palma Bay, an increasing in-migrant population within the Project area, training and employment of local people and use of local goods and services.

Figure 11 Dependence on Natural Resource



Impacts due to physical and economic displacement:

The Project expects that everyone residing within the Afungi Project Site will be permanently displaced resulting in loss of dwellings and infrastructure associated with the household, livelihood activities and community assets within the Afungi Project Site. Access to areas of collective natural resource value (eg forests, wooded grassland, flood plains/ lowlands, dune shrub vegetation, fruit trees, and coconut plantations) will be permanently lost due to the acquisition of land required for the Project. In addition, fishing and sea-based transportation will be affected by Project activities in Palma Bay during the construction and operational phases of the Project due to increased vessel traffic and safety exclusion zones around Project infrastructure. The impacts associated with physical and economic displacement are expected to be of MAJOR significance both during the construction and operational phases. Following the implementation of the Resettlement Action Plan (RAP), the significance will be reduced to MODERATE.





Impacts on tourism: The tourism establishments situated on the islands of Tecomaji, Rongui and Queramimbi will be affected by visual and noise impacts resulting from Project construction activities as well as the presence of the Project and vessel traffic during the operational phase. The disruption and loss of attraction to tourism destinations is expected to be an impact of MAJOR significance. This will be reduced to MODERATE with the implementation of mitigation measures including the design of facilities to minimise visual intrusion on tourism receptors and through appropriate ongoing engagement with tourism operators. While the Project activities are anticipated to negatively influence tourism, there will be a positive impact associated with potential increase in use of tourist facilities by the Project. A MODERATE positive impact is expected as a result of increasing demand for accommodation and use of resorts for leisure purposes during both the construction and operational phases.

Impacts due to in-migration: A wide range of direct and indirect negative environmental, social, and economic impacts on local receptors in Project's host communities such as social services, infrastructure and utilities, social dynamics and cultural life, economy and livelihoods and community health are associated with Project related in-migration, leading to a potential deterioration in the socio-economic environment of the Project's host communities. These impacts are likely to be MAJOR. With appropriate mitigation, including engagement and coordination with relevant authorities impact significance will be reduced to MODERATE.

Impacts on community health: Community health is likely to experience MAJOR significant impacts from the presence of the Project workforce, and due to Project related in-migration into the wider area as well such as an increase in communicable diseases (eg respiratory disease and sexually-transmitted infections and vector borne diseases such as malaria). Similarly, MAJOR significant impacts to community health may arise indirectly from Project activities or associated with Project related in-migration (such as increased pressure through demand on health infrastructure, food and nutrition related issues, community accidents and injuries and soil, water and waste borne diseases). With the implementation of mitigation measures, most impacts are reduced to MODERATE significance, and some MINOR. During the construction and operational phase, the potential for increased sexually-transmitted infections and high risk sexual practices may remain as MODERATE to MAJOR significance post-mitigation.

Enhancement measures to be implemented will promote community health benefits from the presence of the Project. MODERATE to MAJOR positive impacts on community health are anticipated particularly during the operational phase of the Project.

Impacts on local, regional and national economy: The Project will have a range of positive economic impacts at the local, regional, and national levels, including income growth (linked to employment and procurement opportunities), capacity development and increased government revenue. These will range from MINOR positive to MAJOR positive in significance and occur during both the construction and operational phases of the Project. The Project will implement measures to enhance these economic benefits.

There are high expectations associated with the Project to provide opportunities for employment and procurement of goods and services locally and regionally and general economic development. A potential negative impact is associated with unmet community expectations in relation to these opportunities and impact significance is expected to be MODERATE during all Project phases, even with the implementation of mitigation measures.

Impacts due to increase in marine vessel movement: Impacts to international maritime traffic and national and regional cabotage will range from NEGLIGIBLE to MODERATE significance due to the presence of Project vessels and the designation of exclusions zones around Project infrastructure, construction areas and vessels. All impacts will be reduced to NEGLIGIBLE to MINOR once mitigation measures are implemented.

Commercial fishing activities are expected to experience impacts of MINOR to NEGLIGIBLE significance. These potential impacts will be confined to the construction phase of the Project and are most likely to occur in the deep water offshore. Temporary exclusions zones in the offshore gas field and pipeline corridor may impede access to commercial fishing grounds. Impacts are expected to be NEGIGIBLE post-mitigation.

Impacts on archaeology: Impacts to onshore archaeology and cultural heritage are expected to be of MODERATE significance during site clearance and construction activities. However impacts will be reduced to MINOR significance with the implementation of the proposed mitigation. During the operational phase impacts will be NEGLIGIBLE.

Impacts to offshore archaeology and cultural heritage are expected to be MINOR during the construction and operational phases prior to mitigation and NEGLIGIBLE following the implementation of the mitigation measures. All other impacts to mammals will be reduced to MINOR post-mitigation.



Summary of Unplanned Events

An unplanned event is defined as 'a reasonably foreseeable event' that is not planned to occur as part of the Project, but which may conceivably occur as a result of Project activities (eg accidents), even with a low probability. Unplanned events can occur at any phase of the Project onshore and offshore.

The consideration of unplanned events in the EIA has focused primarily on the risks of:

- large volume spills of chemicals or hydrocarbons; and
- failure of subsea infrastructure (such as well blowout or pipeline failure).

Although unlikely to occur should large volumes of hydrocarbon spill in the marine environment the potential consequence to biophysical and socioeconomic receptors could be MAJOR. Sensitive habitats (such as corals and mangroves), local communities dependent on fisheries, as well as tourism operators or aquaculture farms in the vicinity would likely be adversely impacted by such an unplanned event.

Impacts associated with the release of large volumes of natural gas in the marine environment (from failure to or damage of subsea infrastructure) will be less significant.

During the FEED process the Project will identify, assess, mitigate and manage health and safety hazards and risks associated with the Project through a Safety Case. The findings of which will inform decisions related to the design and layout of the Project to reduce the likelihood of unplanned events occurring.

The Emergency Response Plan for the Project sets out response and preparedness measures and incorporates an Oil Spill Contingency Plan. These measures serve to reduce the likelihood, extent and duration of adverse impacts if an unplanned event occurs and establish effective response mechanisms to minimize impacts should such an unlikely event occur.

Cumulative Impacts

It is likely that a number of Mozambique's offshore natural gas resources will be exploited for gas in the coming years resulting in rapid growth in Cabo Delgado Province as a result of the hydrocarbon industry. The cumulative impacts in the region, both positive and negative, as a result are likely to be significant over the life of the Project and beyond. The following have been considered as 'reasonably defined' future developments that could act together with the Project to cumulatively affect the environment:

- establishment of an Industrial Zone (IDZ) by the Government of Mozambique in the vicinity of, or incorporating, the Afungi Project Site.
- future phases of exploration and development of hydrocarbon resources by AMA1, eni and others.

Each of these developments alone has the potential to cause positive and negative biophysical and socio-economic impacts. Cumulatively, the developments will result in significant economic development of the Cabo Delgado Province and the country. The overarching negative cumulative impact of the industrialisation of this area will be the loss of some natural resources as the underdeveloped areas are transformed.

The establishment of the IDZ may result in direct and indirect environmental and social impacts at the local level, but in the long term, consolidating the Project and other operators within the IDZ will serve to confine impacts to one area and ideally allow for easier management of such impacts. Therefore, if managed properly, the IDZ could have an overall positive impact on both biophysical resources and on socio-economic receptors at the regional level given that the Mozambican hydrocarbon resources in this region will undoubtedly be developed. Strategic spatial planning by the Government of Mozambique and its agencies at this early stage is important to promote sustainable development in the region.



ESMP

Environmental and Social Management Plan (ESMP)

A number of measures to manage residual impacts are captured in the Project Environmental and Social Management Plan (ESMP) and other specific management plans have been developed according to the requirements of Mozambican law and good international industry practice. The ESMP and other plans (listed below) will be implemented during construction and operation of the project.

Environmental Education and Training: This is included in Chapter 17 of the EIA Report and describes how Project staff will be made aware of the environmental and social management measures and controls that they will be required to implement.

Project ESMP: This is a tabulated list of environmental and social management measures and is included in Annex D of the EIA Report. The ESMP references the impacts identified in the impact assessment chapters and categorises the management measures according to phase in which they will be implemented and by responsibility for implementation. Specific monitoring requirements have been identified where required.

Waste Management Plan: The Waste Management Plan is included in Annex E of the EIA Report. It defines waste streams, estimates volumes, categorises wastes streams and describes the means by which the Project will manage waste. The Project has committed to following the waste hierarchy:

Reduce the amount of waste generated;

Re-use materials, where appropriate;

Recycle wastes, where appropriate;

Recover as many materials as practical from the remaining waste;

Treat wastes as necessary to render them less hazardous and/or to enable them to be reused or recycled; and

Dispose of residual wastes responsibly at authorised facilities

Decommissioning and Rehabilitation Plan: This plan outlines the Project's commitments to progressively rehabilitate and re-vegetate disturbed areas wherever possible. In line with good international industry practice and the requirements of Mozambican law, this plan provides the principles that will be followed for decommissioning and rehabilitation and commits the Project to reviewing and updating it at least two years before closure. The plan is included in Annex F of the EIA Report.

Emergency Response Plan: This plan is included in Annex H of the EIA Report. It details the emergency organisational structure and protocols that will be implemented to respond to any major incident in a safe, rapid, effective, and efficient manner within the region. Incidents are defined as an event or hazardous circumstance that poses a significant impact to people, the environment, or property.

Initial Resettlement Plan: The Project's Initial Resettlement Plan (IRP) is included in Annex I of the EIA Report. It describes the approach, principles and procedures that will be followed to manage all unavoidable displacement. The IRP provides the framework for the future development of a full Resettlement Action Plan (RAP). The RAP will be prepared in accordance with Mozambican legislation and the International Finance Corporation's Performance Standard 5: Land Acquisition and Involuntary Resettlement (IFC PS 5) in close consultation with the affected communities.



The Draft EIA Report has been finalised in light of all comments received during the public commenting period (27 August to 31 October 2013).

The Final EIA Report has been made submitted to MICOA for decision making (February 2014). If any person or organisation that is interested in or affected by the Project (Interested and Affected Parties (I&APs) has further comments on the Final EIA Report, they are requested to submit these directly to MICOA.





Abbreviations/Units, Acronyms and Glossary

ABBREVIATIONS/UNITS

%	Percent
%PTV	Percentage of pollution-tolerant valves
°C	Degrees Celsius
µg/m³	Micrograms (one-millionth of a gram) per cubic metre
2-D	Two-dimensional
3-D	Three-dimensional
amsl	Above mean sea level
bbl	Barrel
bbl/day	Barrels per day
BBOE	Billion barrels of oil equivalent
BCF	Billion cubic feet
cm	Centimetres
cmol (+)kg-1	Centimoles of positive charge per kilogram
dBA	Decibels (A-weighted scale)
ha	Hectare
hr	Hour
kg	Kilogram
km	Kilometre
km ²	Square kilometres
1	Litre
LAmin	Minimum measured noise level in measurement period
LAeq	Equivalent Continuous Sound Level
LA90	Noise levels that are exceeded for 90% of each sample period
LAmax	Maximum measured noise level in measurement period
l/s	Litres per second
m	Metre
mg	Milligram
m bgl	Metres below ground level
m/s	Metres per second
<u>m²</u>	Square metres
<u>m³</u>	
<u>m³/s</u>	Cubic metres per second
mg.1-1	Milligrams per litre
mi	Millittre
mm	Millimetre
MMtpa	Million metric tons per annum
Nim/yr	Millimetres per year
m5/m	Million metric tene of eacher disuide equivalent
MtCO ₂ e	Million metric tons of carbon dioxide equivalent
MILA	Manuficon motion
	Nozanibicali neucai
IIIII NITLI	Nanonetrie Turbi dity. Unit
nH	Measure of acid or alkaline
pm	Parts per million
ppm	Part per million by volume
ppinv	Parts per thousand
PSU	Practical salinity unit
rme	Root mean squared
	Second
+	Tonne
TCF	Trillion cubic feet
tCO2e	Metric tops of carbon dioxide equivalent
USD	United States dollar
000	onnea oures donai

AC	Agulhas Current
ADI	Area of Direct Influence
ADNAP	Administração Nacional das Pescas (National Fishing Administration)
AFE	Afungi Test Site East
AFT	Afungi Test Site West
Ag	Silver
AII	Area of Indirect Influence
Al	Aluminium
ALARP	As low as reasonably practicable
ALE	Adult literacy and education
AMA1	Anadarko Mocambigue Area 1, Lda
AMODER	Associação Mocambicana para o Desenvolvimento Rural (Mozambican
	Association for Rural Development)
ANC	antenatal care
ANE	Administração Nacional de Estradas (National Roads Administration)
Antarc IW	Antarctic Intermediate Water
APC	Anadarko Petroleum Corporation
	Flementary Polyvalent A gent
	Air Quality Standard
AQ3	All Quality Statuard
AKA	Administração Regional de Águas (Regional Water Administrations)
Area I	Kovuma Basin Offshore Area 1
ARI	Acute respiratory infection
AKI	Antiretroviral treatment
As	Arsenic
ASA	Applied Science Associates Inc
AUV	Autonomous Underwater Vehicle
Ва	Barium
BDI	Biological Diatom Index
Be	Beryllium
BID	Background Information Document
BOD	Biochemical Oxygen Demand
BOP	Blowout Preventer
BTEX	Benzene, toluene, ethyl benzene and xylene (collective name)
C3MR	Propane Pre-cooled Mixed Refrigerant
С	Carbon
Ca	Calcium
CaCO ₃	Calcium carbonate
CBD	Convention on Biological Diversity
CCDA	Complete Community Development Approach
Cd	Cadmium
CEC	Cation exchange capacity
CENACARTA	Centro Nacional de Cartografia e Teledetecção (National Centre for
	Cartography and Tele-detection)
CFM	Caminhos de Ferro de Moçambique EP (Mozambique Ports and Railways
	Company)
CGPCS	International Contact Group on Piracy off the Coast of Somalia
Ch	Champagne soil type
CH ₄	Methane
CHA	Community health agent
CSW	Commercial sex worker
CITES	Convention for the Trade in Endangered Species
Cl	Chloride
CLO	Community Liaison Officer
Со	Cobalt
СО	Carbon monoxide

	Chamical Organ Damand
COLDECa	Chemical Oxygen Demand
COLKEGS	Convention on the International Regulations for the Prevention of Collision
CORDIO	at Sea
CORDIO	Coral Reef Degradation in the Indian Ocean
cor	Crown-ot-thorns (starfish)
Cr	Chromium
CS	Cutter Suction (Dredger)
CSW	Commercial sex workers
CTD	Conductivity-temperature-depth
Cu	Copper
CW	(Indian Ocean) Central Water
DBH	Diameter at breast height
DD	Data Deficient
DEAS	District Economic Activities Service
DFMQAMRP	Darwin/Frontier Mozambique Quirimbas Archipelago Marine Research
	Programme
DINAGECA	National Department of Geography and Cadastre
DMR	Dual Mixed Refrigerant
DNAIA	Direcção Nacional de Avaliação do Impacto Ambiental (National Directora
210111	of Environmental Impact Assessment)
DO	Discolved ovvgen
DOTS	Dissolved oxygen Directly Observed Treatment Short Course
	Duramically positioned
	Dynamicany positioned
DPCA	Direcções Provinciais para Coordenação da Acção Ambiental (Provinciai
DDD	Directorate for Coordination of Environmental Affairs)
DRP	Decommissioning and Rehabilitation Plan
DUAT	Direito de Uso e Aproveitamento da Terra (Land Use and Benefit Rights)
DWAF	South African Department of Water Affairs and Forestry
EACC	East African Coastal Current
EAME	Eastern Africa Marine Ecoregion
EBI	Energy and Biodiversity Initiative
EC	Electrical conductivity
ECAQS	European Commission Air Quality Standard
ECAs	Export Credit Agencies
EEZ	Exclusive Economic Zone
EGPAF	Elizabeth Glaser Paediatric AIDS Foundation
EHA	Environmental health area
EHS	Environmental, Health and Safety
EIA	Environmental Impact Assessment
EIR	Environmental Impact Report
EITI	Extractive Industry Transparency Initiative
EMC	Fast Madagascar Current
FMODRACA	Empresa Mocambicana de Dragagem (Mozambican Drodging Company)
EMDINAGA	Empresa Woçanibicana de Diagageni (Wozanibican Dieuging Company)
ENI	Environmental Management Custors
ENIL	Environmental Management System
ENH	Empresa Nacional de Hidrocarbonetos (National Hydrocarbon Company)
eni	Eni East Africa S.p.A
ENSO	El Nino Southern Oscillation
EP1	First Degree of Primary Education
EP2	Second Degree of Primary Education
EPC	Engineering, Procurement and Construction
EPCC	Exploration and Production Concession Contract
EPDA	Estudo de Pré-viabilidade e Definição do Âmbito (Environmental Pre-
	feasibility Report and Scope Definition)
	Ephemeroptera, Plecoptera and Trichoptera
EPT	
EPT EqIOW	Equatorial Indian Ocean Tropical Surface Water
EPT EqIOW EOT	Equatorial Indian Ocean Tropical Surface Water Environmental Quality Target
EPT EqIOW EQT ERM	Equatorial Indian Ocean Tropical Surface Water Environmental Quality Target Environmental Resources Management Southern Africa (Ptv) Ltd

ESD	Equivalent spherical diameter
ESD/D	Emergency Shutdown and Detection
ESG1	First Cycle of Secondary Education
ESG2	Second Cycle of Secondary Education
ESMP	Environmental and Social Management Plan
ESMS	Environmental and Social Management System (ESMS)
ESP	Exchangeable sodium percentage
FT	Evapotranspiration
FTB	Basic Technical Education
ETE	Elementary Technical Education
ETL	Intermediate Technical Education
EU	
re	
FEED	Front-end Engineering and Design
FEWS NET	Famine Early Warning System Network
FFB	Far-field site
FFP	Fundo de Fomento Pesqueiro (Fisheries Development Fund)
FGD	Focus group discussion
FH	Food for the Hungry
FIPAG	Fundo de Investimento e Património do Abastecimento de Água (The
	Mozambique Water Authority)
FPIC	Free, prior and informed consent
FPU	Floating Production Unit
FTU	Formazin turbidity units
GAPI	Instituição Financeira de Desenvolvimento de Moçambique (Mozambican
	Institute of Development Finance)
GDACE	Gauteng Department of Agriculture Conservation and Environment (South
	Africa)
GBDV	Gender-based domestic violence
GDEM	Global Digital Elevation Model
GDP	Gross domestic product
GFF	Global Environment Facility
CHC	Greenhouse ass
CIICNI	The International Crown of Liquefied Natural Cas Importants
CIS	Coographic Information System
CMDSS	Clobal Maritima Distrace and Safaty System
GND55	Giobal Maritime Distress and Safety System
GNP	Gross national product
GOM	Government of Mozambique
GPS	Geographically Positioned System
GWP	Global warming potential
H ₂ O	Water
H_2S	Hydrogen sulphide
HAT	Highest Astronomical Tide
HCO ₃	Bicarbonate
HF	High flow
HFO	Heavy fuel oil
Hg	Mercury
HH	Households
HHV	Higher Heating Value
HIA	Health Impact Assessment
HIV/AIDS	Human immunodeficiency virus/acquired immunodeficiency syndrome
HLV	Heavy Lift Vessel
HSB	Health seeking behaviour
HSS	Health systems strengthening
НУСОМ	HYbrid Coordinate Ocean Model
I&APs	Interested and affected parties
IACM	Instituto de Aviação Civil de Mocambique (National Aviation Institute)
IADC	International Association for Drilling Contractors
IBAc	Important Bird Areas
	International Council on Monuments and Sites
ICOMO5	International Council on Monuments and Siles

IDPPE	Instituto Nacional de Desenvolvimento da Pesca de Pequena Escala (National		
	Small-Scale Fishing Development Institute)		
IDZ	Industrial Development Zone		
IEC	Information, education and communication		
IFC	International Finance Corporation		
ш	Index of Habitat Integrity		
	Index of Habitat Integrity		
	Instituto de Investigação i esqueira (Fisheries Research Institution)		
IMF	International Monetary Fund		
IMO	International Maritime Organisation		
Impacto	Impacto, Projectos e Estudos Ambientais Lda		
INAHINA	Instituto Nacional de Hidrografia e Navegação (National Hydrography and		
	Navigational Institute)		
INAM	Instituto Nacional de Meteorologia (National Institute of Meteorology)		
INAMAR	Instituto Nacional da Marinha (National Maritime Institute)		
INAR	Mozambican National Institute for the Support of Refugees		
INGC	Instituto Nacional de Gestão de Calamidades (National Institute for of		
	Disaster Management)		
INMARSAT	Convention on the International Maritime Satellite Organisation		
INP	Instituto Nacional do Petróleo (National Petroleum Institute)		
IOC	Indian Ocean Commission		
IPCC	Intergevernmental Panel on Climate Change		
	Corporate income fax		
IRPS	Individual Income Tax		
ISO	International Organization for Standardisation		
ITCZ	Intertropical Convergence Zone		
ITN	Insecticide-treated net		
IUCN	International Union for Conservation of Nature		
IWANQ	Mozambique Water Quality Standards		
JAM	Joint Aid Management		
К	Potassium		
Ка	Katspruit soil type		
Kd	Kroonstad soil type		
KII	Key informant interview		
KPI	Key Performance Indicator		
L&FS	Life and Fire Safety		
LAT	Lowest Astronomical Tide		
	Least Concern		
	Light Construction Vessel		
LEV	Light Construction vessel		
LEMO	League of Scouts of Mozambique		
LF	Low flow		
Li	Lithium		
LIA	Late Iron Age		
LNG	Liquefied Natural Gas		
Lo	Kroonstad soil type		
LOLE	Lei dos Órgãos Locais do Estado (Law for Local State Bodies)		
LP	Liquefied petroleum		
LULUCF	Land use, land use change and forestry		
LZ	Lower Zone		
MARPOL 73/78	International Convention for the Protection of Pollution from Ships		
MBREMP	Mnazi Bay-Royuma Estuary Marine Park		
MC	Mozambique Current		
MCA	Millonnium Challongo Account		
MCE	Marambiana Channal Eddias		
MDEA			
MDEA			
MdP	Mocimboa da Praia		
MEG	Monoethylene glycol		
Mg	magnesium		
MHWN	Mean High Water Neap		
MHWS	Mean High Water Spring		

MICOA	Ministério para a Coordenação da AcçãoAmbiental (Ministry for the
	Coordination of Environmental Affairs)
MINAG	Ministério da Agricultura (Ministry of Agriculture)
MIRAI	Macro-invertebrate Response Index
MIREM	Ministério dos Recursos Minerais (Ministry of Mineral Resources)
MISAU	Ministério da Saúde (Ministry of Health)
MITUR	Ministério do Turismo (Ministry of Tourism)
MLWN	Mean Low Water Neap
MLWS	Mean Low Water Spring
MMO	Marine mammal observer
Mn	Manganese
Mo	Malybdonum
MOE	Motovial Offloading Facility
MDA	Material Onloading Facility
MDD	Multinum and Deale
MPD	Multipurpose Dock
MPescas	Ministerio das Pescas (Ministry of Fisning)
MR	Mixed refrigeration
MSI	Metocean Services International (Pty) Ltd
MSL	Mean sea level
MZ	Marginal Zone
Na	Sodium
NABM	Non-aqueous based muds
NaCl	Sodium chloride
NADW	North Atlantic Deep Water
Nb	Namib soil type
NCD	Non-communicable disease
NCEP	National Centre for Environmental Predictions
NEMC	North East Madagascar Current
NEMBA	National Environmental Management: Biodiversity Act (No. 20 of 2004)
NFB	Near-field site
NGO	Non-governmental organisation
NH4	Ammonium
Ni	Nickel
NIDW	North Indian Dean Water
NMCP	National Malaria Control Programme
NOAA	National Malaria Control Programme
NOAA	National Oceanic and Atmospheric Administration
NO ₂	
NO ₃	Nitrate
NOx	Nitrogen oxides
NORM	Naturally occurring radioactive materials
NSR	Noise sensitive receptor
NT	Near Threatened
NTS	Non-technical Summary
NTU	Nephelometric Turbidity Unit
NW	North-west
OBM	oil-based muds
OGP	International Oil and Gas Producers Association
OILPOL	International Convention for the Prevention of Pollution of the Sea by Oil
OPRC	Oil Pollution Preparedness, Response and Cooperation Convention
OSCP	Oil Spill Contingency Plan
Р	Phosphorus
PAC	Potentially affected communities
PAH	Polyaromatic hydrocarbon
PARP	Poverty Reduction Action Plan
PATI	Priority area for tourism investment
Pb	Lead
PC	Project Contribution
PCMRB	Petronas Carigali Mozambique Rovuma Basin
PCSW	Potentially contaminated surface water
1 (377	
PEC	Productod Environmontal Concentration

PES	Present ecological state	
PESPA	Plano Estratégico do Subsector da Pesca Artesanal (Strategic Plan for the	
	Artisanal Fisheries Subsector)	
PGU	Power Generation Unit	
PICT	Provider initiated counselling and testing	
Pig	Pipeline integrity gauge	
PIIM	Project Induced In-Migration	
PLFT	Pipeline End Termination Structure	
PLONAR	Pose Little or No Biele to the Environment	
DM	Posticulate matter	
DM10	Particulate matter < 10	
PMIU	Particulate matter < 10µm	
PMI		
PMICI	Prevention of mother-to-child transmission	
PMW	Palma West Site	
PNA	National Water Policy	
POM	Particulate organic matter	
PPE	Personal protective equipment	
PR	Propane Refrigeration	
PSU	Practical salinity units	
PTS	Permanent Threshold Shift	
PTV	Pollution-tolerant valves	
PTW	Permit to work	
PPR	Public Participation Report	
RAP	Resettlement Action Plan	
RDT	Rapid diagnostic test	
RHIA	Rapid Health Impact Assessment	
RHMIS	Routing Health Information Management System	
REINIS	Routine Treatilit information Management System	
RO	Reverse osmosis	
RUV	Remotely operated vehicle	
KFF DTA	Resettlement Policy Framework	
RIA	Road traffic accident	
SADC	Southern African Development Community	
SAMSA	South African Maritime Safety Authority	
SAR	IMO International Convention on Search and Rescue	
SASS5	South African Scoring System, version 5	
SAWQG	South African Water Quality Guidelines	
Sb	Antimony	
SBM	Synthetic-based muds	
SCS (Method)	Storage Conservation Service (Method)	
SDoH	Social determinants of health	
Se	Selenium	
SE	South-east	
SEC	South Equatorial Current	
SEL	Sound exposure level	
SEMC	South East Madagascar Current	
SEP	Stakeholder Engagement Plan	
SES	Simplified Environmental Study	
SI	Surface irradiance	
SMA	Soil Moisture Accounting	
SMA	Single Mixed Petrigerent	
Sivin		
Sn		
502		
<u>504</u>	Sulphate	
SOLAS 74	International Convention for the Safety of Life at Sea	
SPI	Specific Pollution Sensitivity Index	
SPS	Subsea Production System	
Sr	Strontium	
SS	Suspended solids	
SSS	Side Scan Sonar	

STCW	IMO International Convention on Standards of Training, Certification and
	Watch keeping for Seafarers
STH	Soil-transmitted helminth
STI	Sexually transmitted infection
Subt SIOW	Indian Ocean Subtropical Surface Water
S-Unit	Sand unit
TB	Tuberculosis
Tc	Travel time
TCN	Third country national
TDS	Total Dissolved Solids
TH	Traditional healer
ToR	Terms of Reference
TS	temperature/salinity
TSHD	Trailer Suction Hopper Dredger
TSS	Total Suspended Solids
TTS	Temporary Threshold Shift
TWC	Temporary workers camp
TWOR	Target Water Quality Range
	Uranium
UEM	Universidade Eduardo Mondlane (Eduardo Mondlane University)
UNI	United Nations
UNCLOS	UN Law of the Sea Convention
UNCLOS	United Nations Environment Programme
UNESCO	United Nations Educational Scientific and Cultural Organization
UNECCC	United Nations Educational, Scientific and Cultural Organization
UNFCCC	United Nations Mater Quality Cuidalines
UNWQG	United Nations water Quality Guidelines
	United States of America
USAID	United States Agency for International Development
USEFA	Ultraviolet
	Unaverladed Ordinance
	Unper Zene
V	Vanadium
VAI	Value added tax
VEGRAI	Vegetation Response Assessment Index
VOC	Volatile Organic Compounds
VU	Vulnerable
WaMP	Water Management Plan
WB	World Bank
WBM	Water-based muds
WHO	World Health Organization
WIO	West Indian Ocean
WIOLAB	West Indian Ocean Land Based
WMP	Waste Management Plan
WMS	Watershed Modelling System
WQ	Water quality
W-Unit	Wetland unit
WWF	Worldwide Fund for Nature
Zn	Zinc
ZSL	Zoological Society of London
ZTV	Zone of Theoretical Visibility

GLOSSARY

Afungi Project Site	The location on Afungi Peninsula where the LNG Facility and associated
	intrastructure (roads, airstrip, accommodation, etc) will be located. This is the area where the Project has acquired the rights to use the land.
Afungi Project Site	This refers to Afungi Project Site and the areas around the Afungi Project
and Surrounds	Site where people's access to livelihood assets will be affected.
Afungi Site	The Site Improvement activities are intended to upgrade existing
Improvement	infrastructure and make minor preparations to facilitate the construction
activities	phase, should the Project be approved.
Alternative	Alternatives can refer to any of the following, but are not limited to:
	alternative sites for development, alternative projects for a particular site,
	alternative site layouts, alternative designs, alternative processes and
	alternative materials.
Anthropogenic	Caused by humans.
Aquifer	Fresh water (usually) held in underground layers of water-bearing
-	permeable rock or unconsolidated sediments (gravels/sand/silt).
Arable	Land that can be cultivated for growing crops.
Area 1	Area 1 is the area licensed to AMA1 by the Ministry of Energy and
	Mineral Resources for the exploration and development of
	hydrocarbons.
Area 1 Gas Fields	The offshore AMA1 gas fields, namely Golfinho and Prosperidade,
	found in Area 1.
Area 4	Area 4 is the area licensed to eni by the Ministry of Energy and Mineral
	Resources for the exploration and development of hydrocarbons.
Area 4 Gas Fields	The offshore eni gas field covered in the EIA, namely Mamba, found in
	Area 4.
Area of Direct	The ADI comprises the areas offshore, near shore and onshore where the
Influence (ADI)	Project infrastructure will be located, ie the direct Project Footprint. The
	ADI comprises two broadly defined areas of interest:
	Offshore Project Footprint Area; and
	Offshore Project Footprint Area; andAfungi Project Site and Surrounds.
Area of Indirect	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises:
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises:
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas;
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas;
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted;
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted;
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The All refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The All refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly,
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale.
Area of Indirect Influence (AII)	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale.
Area of Indirect Influence (AII) Avifauna Bairros	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale. Birds.
Area of Indirect Influence (AII) Avifauna Bairros Baseline	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale. Birds.
Area of Indirect Influence (AII) Avifauna Bairros Baseline	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale. Birds. Neighbourhoods. The current physical, biological, cultural and human conditions that will prevail in the absence of the Project, including interactions among them.
Area of Indirect Influence (AII) Avifauna Bairros Baseline Bathymetry	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale. Birds. Neighbourhoods. The current physical, biological, cultural and human conditions that will prevail in the absence of the Project, including interactions among them.
Area of Indirect Influence (AII) Avifauna Bairros Baseline Bathymetry	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale. Birds. Neighbourhoods. The current physical, biological, cultural and human conditions that will prevail in the absence of the Project, including interactions among them. Underwater depths. Bathymetry maps show underwater depths at various locations.
Area of Indirect Influence (AII) Avifauna Bairros Baseline Bathymetry Benthic zone	 Offshore Project Footprint Area; and Afungi Project Site and Surrounds. The AII refers to areas where the Project Footprint has an indirect influence, and comprises: Palma Bay and offshore deepwater in the vicinity of the production areas; Palma Town and villages or settlements outside the Afungi Project Site that are indirectly impacted; the islands at the mouth of Palma Bay: Tecomaji, Rongui and Queramimbi islands, and possibly islands further south along the coast eg Vamizi Island; and the broader district, province and potentially region can also be included in the AII from a socio-economic perspective. Similarly, the AII potentially extends to a national level, as the Project may have benefits at a national scale. Birds. Neighbourhoods. The current physical, biological, cultural and human conditions that will prevail in the absence of the Project, including interactions among them. Underwater depths. Bathymetry maps show underwater depths at various locations.

Benthos	Species living in or on ocean, sea, lake beds and/or river bottoms.
Berthing Area	The area of the Near Shore infrastructure where LNG Carriers will berth.
Biodiversity	The variability among living organisms – animals, plants, their habitats –
	from all sources including terrestrial, marine and other aquatic
	ecosystems, and the ecological complexes of which they are part. This
	includes diversity within species, between species, and of ecosystems.
Biomass	The total mass of living matter within a given unit of environmental area.
Biome	Major ecological community, a division of the world's vegetation that
	corresponds to a particular climate and is characterised by certain types
	of plants and animals.
Biota	The animal and plant life of a given region.
Biotope	An area of uniform environmental conditions providing a living place
	for a specific assemblage of plants and animals. Biotope is almost
	synonymous with the term habitat, but while the subject of a habitat is a
	species or a population, the subject of a biotope is an interdependent
	biological community.
Bioturbation	The displacement and mixing of sediment particles by fauna or flora.
Brine	Concentrated salt water from the Desalination Plant.
Bund	Containment wall to capture spills before loss to the environment.
Carbon dioxide	A metric measure used to compare the emissions from various
equivalent	greenhouse gases based upon their global warming potential (GWP).
Casing	A process carried out to isolate portions of a well so as to protect the
	aquifers of groundwater, as well as to provide a support structure to the
0.1.1	well itself.
Catchment	An area from which surface run-off is carried away by a single drainage
Calaasa	system.
Cetacea	Order of marine mammals containing dolphins, whales and porpoises.
Chlorophyll	A substance used to retard a chemical reaction.
Chorophyn Cloan surfaco run	Bun off from areas doomed not at risk to oil or chemical contamination
off	Clean surface run off will be allowed to discharge to the surrounding
011	environment
Condensate	Hydrocarbon liquid that condenses from natural gas when the pressure
condensate	drops below dewpoint.
Coral bommies	A coral bommie (from 'bombora' – an Aboriginal word meaning outcrop
corta commutes	or mountain or reef) is either an outcrop of coral reef, often resembling a
	column that is higher than the surrounding platform of reef, or an
	isolated piece of reef on sand.
Coral fringing reef	Coral reef on outside of shores or islands.
Coral rubble	Unconsolidated fragments of broken dead coral
Corals	Marine invertebrates in the class Anthoroa of the phylum Chidaria
Cordis	typically living in compact colonies of many identical individual polyns
	The group includes the important reef builders that inhabit tropical
	oceans, which secrete $CaCO_3$ to form a hard skeleton.
Cruchagoang	Arthurned out has make and proving that have joined appendance and
Clustacealls	hard outor cholle
Culvert	A man-made structure used to channel water
Curvert Cumulative impacts	Impacts that result from the incremental impact of the proposed activity
Cumulative impacts	on areas or common resources used or directly impacted by the Project
	from other existing planned or reasonably defined dovelopments at the
	time the risks and impacts identification process is conducted when
	added to the impacts of other past present or reasonably foreseeable
	future activities Cumulative impacts can occur from the collective
	impacts of individual minor actions over a period of time and can
	include both direct and indirect impacts
Cuttings	Debris (pieces of sand, gravel and rock) that are brought to the surface
Cattings	from the well during drilling
	nom ale wen dunitg annung.

Demersal Demersal zone Dewatering Dhows Diatoms Direct impacts	not dependent on surface rainfall run-off, but rather water welling up from below. <i>Dambos</i> are also found at higher elevations outside wetland systems and are presumably fed from arterial springs, and retain water for longer periods during the dry season. The part of the water column that is near to the seabed and the benthos. Demersal fish are those that spend the majority of their life cycle on or near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal Demersal zone Dewatering Dhows Diatoms Direct impacts	from below. <i>Dambos</i> are also found at higher elevations outside wetland systems and are presumably fed from arterial springs, and retain water for longer periods during the dry season. The part of the water column that is near to the seabed and the benthos. Demersal fish are those that spend the majority of their life cycle on or near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal Demersal zone Dewatering Dhows Diatoms Direct impacts	systems and are presumably fed from arterial springs, and retain water for longer periods during the dry season. The part of the water column that is near to the seabed and the benthos. Demersal fish are those that spend the majority of their life cycle on or near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal Demersal zone Dewatering Dhows Diatoms Direct impacts	for longer periods during the dry season. The part of the water column that is near to the seabed and the benthos. Demersal fish are those that spend the majority of their life cycle on or near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal Demersal zone Dewatering Dhows Diatoms Direct impacts	The part of the water column that is near to the seabed and the benthos. Demersal fish are those that spend the majority of their life cycle on or near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal zone Dewatering Dhows Diatoms Direct impacts	Demersal fish are those that spend the majority of their life cycle on or near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal zone Dewatering Dhows Diatoms Direct impacts	near to the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Demersal zone Dewatering Dhows Diatoms Direct impacts	That part of the seabed. That part of the sea or ocean or deep lake comprising the lower water column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Dewatering Dows Diatoms Direct impacts	Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Dewatering Dhows Diatoms Direct impacts	Column that is near to and affected by the sea/lake bed and benthos. Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Dewatering Dhows Diatoms Direct impacts	Removal of water from. A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Dhows Diatoms Direct impacts	A lateen-rigged ship with one or two masts, used in the Indian Ocean. A single-celled algae that has a cell wall of silica.
Diatoms Direct impacts	A single-celled algae that has a cell wall of silica.
Direct impacts	0 0
	Impacts that result from a direct interaction between a planned Project
	activity and the receiving environment/receptors (eg between
	occupation of a site and the pre-existing habitats, or between an effluent
	discharge and receiving water quality) These impacts are usually
	associated with the construction operation or maintenance of an activity
	associated with the construction, operation of maintenance of an activity,
Domostia cohoto co	Transmost between two noints within a sountry
	The second the second s
Jrilling fluids	These are used to control subsurface pressures, lubricate the drill bit,
drilling muds)	stabilise the well bore and carry the cuttings to the surface.
OUAT process	The DUAT or <i>direito de uso e aproveitamento da terra</i> process refers to the
	process of obtaining right of use of land in Mozambique.
Jynamically	The location or position of the vessel is maintained by the vessel's
positioned	specialised propulsion/station-keeping system.
Early works	These works will involve site preparation activities and will expand on
	the activities started in the Afungi Site Improvement phase, and are
	anticipated to begin following MICOA approval of this EIA.
Ecosystem	A community of plants, animals and smaller organisms that live, feed,
5	reproduce and interact in the same area or environment.
Ecosystem services	The benefits people obtain from ecosystems.
EIA Pre-assessment	This phase involves submitting a Pre-assessment Application to the
Application Phase	Provincial Directorate of Environmental Affairs (Direcção Provincial de
Screening)	Coordenação de Acção Ambiental or $DPCA$) which then categorises the
Screening)	Project based on the level of environmental assessment required
7	Plan to a draw contingencies accordent with an eccord whether d
Imergency	Plan to address contingencies associated with process upset and
Response Plan	accidental circumstances.
Indemic	Restricted to a particular area. Used to describe a species or organism
	that is confined to a particular geographical region.
Engineering Team	The Project's Engineering Team (involves FEED Contractors involved in
	the design of the Project).
Environment	The surroundings within which humans exist, and that are made up of:
	(i) the land, water and atmosphere of the earth;
	(ii) micro-organisms, plant and animal life:
	(ii) micro organismo, plant and animar me,
	(iii) any part or combination of (i) and (ii) and the inter-relationships
	among and between them; and
	among and between ment, and
	(iv) the physical, chemical, aesthetic and cultural properties and
	conditions of the foregoing that influence human health and well-
	being. This includes the economic, social, cultural, historical and
	political circumstances conditions and objects that affect the
	romitian encambrances, continuous and objects that affect the
	existence and development of an individual, organism or group.
Environmental and	existence and development of an individual, organism or group. A set of actions and measures to be carried out by the proponent, in
Environmental and Social Management	A set of actions and measures to be carried out by the proponent, in order to manage the negative impacts and maximise the positive impacts
Environmental and Social Management Pan (ESMP)	A set of actions and measures to be carried out by the proponent, in order to manage the negative impacts and maximise the positive impacts resulting from the implementation of the proposed activity, prepared

Environmental	The process of predicting and evaluating the social and environmental
Impact Assessment	impacts and risks of a proposed Project, and identifying mitigation
	measures that will enable the Project to meet the requirements of
	stakeholders, applicable laws and regulations, and any additional
	requirements for social or environmental performance identified by the
	Project, and so that impacts are as low as technically and financially
	feasible.
Environmental	Ensuring that environmental concerns are included in all stages of
management	development, so that development is sustainable and does not exceed the
	carrying capacity of the environment.
Epifauna	Animals that live on the surface of the substrata.
Estuaries	A partly enclosed body of water that has one or more rivers flowing into
	it and is connected to the open sea. Estuaries thus receive fresh water
	from inland and salt water from the sea. Tides influence how much (and
	how far) salt water intrudes into the estuary and river.
Estudo de Pré-	The scoping phase or EPDA phase, as it is known in Mozambique, aims
Viabilidade	to identify key issues and concerns associated with the proposed
Ambiental e	development. These could include Project-related activities that may
Definição de	have the potential to contribute to or cause potentially significant
Âmbito (EPDA)	impacts to environmental and socio-economic receptors and resources in
phase or scoping	the area
phase of scoping	
Evolution Zono	Area from which weesals are prohibited from optoring
Exclusive Economic	A concent adopted at the Third United Nations Conference on the Law
Zono (FEZ)	of the See (1982), whereby a coastal State assumes jurisdiction over the
ZOHE (EEZ)	of the Sea (1962), whereby a coastal state assumes jurisdiction over the
	exploration and exploration of marine resources in its adjacent section of
	the continental shell, taken to be a band extending 200 miles from the
Fauna	Collectively, the animal life of a region.
Γ^{*}	Could and the Could and the Could and the Armalian strength and the Could and the Coul
Fishing Centres	Settlements of fishers, where fishers and fish traders live on a permanent
Fishing Centres	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis.
Fishing Centres Flaring	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases.
Fishing Centres Flaring Flora	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region.
Fishing Centres Flaring Flora Formation or	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during
Flaring Flora Formation or produced water	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations.
Flaring Flora Formation or produced water Front-end Engineering Design	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED)	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Eugiting amissions	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GUG)	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ ,
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG)	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO2, CH4, N2O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO2, CH4, N2O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain hydrocarbons.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain hydrocarbons. The filling of equipment or pipelines with water to test for any losses of
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons Hydrotesting	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain hydrocarbons. The filling of equipment or pipelines with water to test for any losses of pressure through leaks.
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons Ichthyofauna	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain hydrocarbons. The filling of equipment or pipelines with water to test for any losses of pressure through leaks. Fish.
Fishing Centres Flaring Flora Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrocarbons Hydrotesting Ichthyofauna Impact	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO2, CH4, N2O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain hydrocarbons. The filling of equipment or pipelines with water to test for any losses of pressure through leaks. Fish. Any alteration of existing conditions. adverse or beneficial. caused
Fishing Centres Flaring Flora Formation or produced water Front-end Engineering Design (FEED) Fugitive emissions Gender diglossia Greenhouse gases (GHG) Habitat Hazardous waste Herpetofauna Hydrotesting Ichthyofauna Impact	Settlements of fishers, where fishers and fish traders live on a permanent or seasonal basis. The safe burning of excess unwanted gases. Collectively, the plant life of a region. Water trapped in the gas and rocks that is brought to the surface during operations. The design phase of the Project. The unintended escape of greenhouse gas emissions. A situation in which two different languages are used by men and women within a community. Unless indicated otherwise, GHG emissions are made up of CO ₂ , CH ₄ , N ₂ O, HFCs, PFCs and SF6. An ecological or environmental area that is inhabited by a particular species of animal, plant or other type of organism. It is the natural environment in which an organism lives, or the physical environment that surrounds (influences and is used by) a species population. Substances classified as hazardous wastes possess at least one of four characteristics of signitability, corrosivity, reactivity or toxicity – or appear on special lists. Amphibians and reptiles. Compounds that contain only hydrogen and carbon molecules. Petroleum-based products such as LNG and condensate contain hydrocarbons. The filling of equipment or pipelines with water to test for any losses of pressure through leaks. Fish. Any alteration of existing conditions, adverse or beneficial, caused directly or indirectly by the Project.

Indirect impacts	Impacts that result from other activities that are encouraged to happen as
-	a consequence of the Project (eg in-migration for employment placing a
	demand on resources). Indirect impacts can also be referred to as
	induced or secondary impacts. These types if impacts include all of the
	potential impacts that do not manifest immediately when the activity is
	undertaken or which occur at a different place as a result of the activity.
Infauna	Animals that live in sediment.
Infectious disease	Illnesses that are attributable to specific infectious agents or their toxic
	products that arise through the transmission of these agents or their
	products from an infected person, animal or inanimate reservoir to a
	susceptible host. Examples include water-borne, water-related, food-
	borne, respiratory and sexually transmitted diseases.
Initial Resettlement	Establishes the principles, approach and procedures that will be followed
Plan	in planning resettlement and awarding compensation to people who are
	physically or economically displaced by a project. The IRP provides the
	framework for the future development of a full Resettlement Action Plan
	(RAP).
Interested and	Individuals, communities or groups, other than the proponent or the
Affected Parties	authorities, whose interests may be positively or negatively affected by
(I&APs)	the proposal or activity and /or who are concerned with a proposal or
(activity and its consequences.
Intertidal zone	The area of the shore that is above water at low tide and under water at
Interneti Zone	high tide
Intertropical	An area of low pressure that forms where the Northeast Trade Winds
Convergence Zone	meet the Southeast Trade Winds near the equator. As these winds
(ITC7)	converge moist air is forced upward causing water vapour to condense
(IICL)	and cool resulting in heavy precipitation
Invasive or alien	Species introduced outside its normal distribution. Its establishment and
species	spread modifies ecosystems habitats or species. Aliens are called
species	invasive when they spread rapidly and outcompete indigenous species
* . 1 .	Control of the species of the second species.
Invertebrates	Species lacking a spinal column (eg crabs)
Invertebrates	Species lacking a spinal column (eg crabs).
IUCN Red List	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological
IUCN Red List	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems
IUCN Red List	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction.
IUCN Red List	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction.
Invertebrates IUCN Red List Keystone species	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which place a gritical rale in maintaining the
Invertebrates IUCN Red List Keystone species	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the
INVERTED IUCN Red List Keystone species	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope.
Invertebrates IUCN Red List Keystone species Land capability	Species lacking a spinal column (eg crabs).Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction.A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope.A combination of terrain form, soil types, thickness and slope gradients
Invertebrates IUCN Red List Keystone species Land capability	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability
Invertebrates IUCN Red List Keystone species Land capability	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area.
Invertebrates IUCN Red List Keystone species Land capability Land cover	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation
Invertebrates IUCN Red List Keystone species Land capability Land cover	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG)	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH ₄) that has been converted to liquid form for ease of storage or transport.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone	Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH ₄) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath).
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty	 Species tacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The jetty infrastructure where LNG Carriers will berth to load LNG for
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty	 Species tacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The jetty infrastructure where LNG Carriers will berth to load LNG for export.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty LNG Facility	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The jetty infrastructure where LNG Carriers will berth to load LNG for export.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty LNG Facility	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The pietty infrastructure where LNG Carriers will berth to load LNG for export. The processing facility where natural gas will be liquefied, and associated facilities.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty LNG Facility LNG Train	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The processing facility where natural gas will be liquefied, and associated facilities. The LNG Plant's liquefaction and purification facilities.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty LNG Facility LNG Train Machambas	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The processing facility where natural gas will be liquefied, and associated facilities. The LNG Plant's liquefaction and purification facilities.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty LNG Facility LNG Train Machambas Macroalgae	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The processing facility where natural gas will be liquefied, and associated facilities. The LNG Plant's liquefaction and purification facilities. Small-scale farms (fragmented cultivated lands). Seaweeds including red, brown and green algae.
Invertebrates IUCN Red List Keystone species Land capability Land cover Landscape Liquefaction Liquefied Natural Gas (LNG) Littoral zone LNG Export Jetty LNG Facility LNG Train Machambas Macroalgae Mangrove	 Species lacking a spinal column (eg crabs). Also known as the IUCN Red Data List, this is the world's most comprehensive inventory of the global conservation status of biological species. Taxa are classified according to formal ranking systems assessing relative risks of extinction. A species that has a disproportionate effect on its environment relative to its abundance, and which plays a critical role in maintaining the structure of an ecological community/biotope. A combination of terrain form, soil types, thickness and slope gradients in association with broad agricultural potential define the land capability of an area. The physical coverage of land, usually expressed in terms of vegetation cover or lack of it. A geographical mosaic composed of interacting ecosystems resulting from the influence of geological, topographical, soil, climatic, biotic and human interactions in a given area. The process by which natural gas is cooled to condense it to a liquid. A natural gas (predominantly methane, CH4) that has been converted to liquid form for ease of storage or transport. That part of the sea that extends from the high water mark to the edge of the continental shelf (commonly defined as the 200m isobath). The processing facility where natural gas will be liquefied, and associated facilities. The LNG Plant's liquefaction and purification facilities. Small-scale farms (fragmented cultivated lands). Seaweeds including red, brown and green algae. Kinds of trees up to medium height, and shrubs, which grow in saline

Marine ornamental fish	Small fish in families commonly used in marine aquaria, including Chaetodontidae, Labridae, Zanclidae, Acanthuridae, Serranidae, Pseudochromidae, Pomacentridae, etc.
Materials Offloading Facility (MOF)	The service port and transit point of equipment and materials used during the construction and operation phase of the Project.
Mesopelagic	The part of the pelagic zone that extends from a depth of 200 to 1,000 below the sea's surface.
Metocean	The combined study of meteorology and oceanography.
Microalgae	Microscopic single-celled algae in fresh, brackish (estuarine) and mar waters. Pelagic species comprise phytoplankton, while those on sediments are benthic microalgae.
Mitigate	The implementation of practical measures to reduce adverse impacts enhance beneficial impacts of an action.
Mitigation measure	A feature, procedure or other action that the Project commits to implement to avoid or reduce the magnitude of an adverse impact, or enhance the magnitude of a positive impact.
Monoethylene glycol (MEG)	A colourless, virtually odourless and slightly viscous liquid used to inhibit the formation of hydrates or condensate by altering the freezin point of the water found in raw natural gas.
Multipurpose Dock (MPD)	Marine structure comprising the MOF, the Subsea Construction Area (SSCA) and Marine Service Facility (MSF).
Nautical mile	A measurement of distance at sea equal to 1,852m.
Near Shore environment	The waters westward of Mozambique's Maritime Baseline.
Near Shore Project	The proposed marine infrastructure within Palma Bay necessary for construction, operation and maintenance of the Project. This includes logistic, support and export facilities (eg shipping channels, MOF and
	LNG Export Jetty).
No-go alternative	The no-go alternative is the option of not undertaking the proposed activity or any of its alternatives. This alternative also provides the baseline against which the impacts of other alternatives should be
Offshore environment	The area beyond Mozambique's Maritime Baseline.
Offshore Project	The proposed offshore production wells and the infrastructure necess
components	to develop the gas reserves. This includes the offshore pipeline system that will convey natural gas from the offshore production field to the onshore facilities.
Offshore Project Footprint Area	The areas in the offshore and near shore environment that may reasonably be expected to be physically touched by Project
	infrastructure, across all phases.
Oligotrophic	Low nutrient conditions.
Onshore environment	The terrestrial environment.
Onshore Project	The proposed onshore LNG facilities and supporting infrastructure (e worker accommodation facilities, construction areas, access roads and airstrip).
Onshore Project	The area onshore that may reasonably be expected to be physically
Footprint Area	touched by Project infrastructure, across all phases.
Operational phase	The period of the Project's life when gas will be extracted, processed a exported.
Oxygen scavengers	Used to reduce the oxygen content in air or water to slow the process corrosion.
Particulate matter	Fine particles of solid matter.
Pelagic	The part of the water column in the open sea that is not closely associated with the seabed.
Pelagic zone	Water body in oceans, seas or lakes that is not close to the bottom or

Pioneer Dock	A Pioneer Dock will be established to allow the import of heavy
Dis alia a insta antita	A pulindrical device incorted into a principal to incore the pine on the
Pipeline integrity	A cylindrical device inserted into a pipeline to inspect the pipe or to
gauge (PIG)	Sweep the pipe clean of water, rust or other foreign matter.
Plankton	inhabit the pelagic zone of water bodies.
Potsherd	A broken piece of ceramic material, especially one found on an
D 1 (1	archaeological site.
Production zones	Agricultural areas.
Project activities	Activities that are planned as part of the Project, including all phases. This excludes unplanned events
Protected areas	An area of land and/or sea especially dedicated to the protection of
1 loteetee ureus	biological diversity, and of natural and associated cultural resources, and
	managed through legal or other effective means.
Public consultation	Term used when referring to the activity of consulting with stakeholders
	or I&APs.
Public participation	Term used when referring to the regulatory-driven stakeholder or I&AP
process	engagement process.
Receptor	Humans and other animals that can be impacted by Project activities.
Regional cabotage	Transport in coastal waters.
Resettlement Policy	Establishes the principles and procedures that will be followed in
Framework	planning resettlement and awarding compensation to people who are
	physically or economically displaced by a project, including the process
	by which people can raise a grievance if they consider that have been
	adversely affected or unfairly treated.
Residual impact	Impacts that remain after mitigation measures, including those
	incorporated into the Project's design, have been applied.
Resource	An element of the physical, biological, cultural or human environment
	that is not a human or other animal (these are referred to as receptors),
	which can be impacted by the Project activities.
Revised Onshore	Following input from various specialists on the sensitivities (biophysical
Project Footprint	and socio-economic) of the Afungi Project Site, a revised site area was
Area	determined, known as the 'Revised Onshore Project Footprint Area '.
	This revised layout was considered in the assessment of the residual
	impact, as the change in layout is considered as a mitigation measure.
Salinity	The dissolved salt content in sea water.
Seagrass	Flowering rooted plants (angiosperms) from one of four plant families all
	in the order <i>Alismatales</i> (in the class of monocotyledons), which grow in
	marine, fully saline environments.
Seascape Unit (S-	Coastal landscape and adjoining areas of open water, including views
Unit)	from land to sea, from sea to land and along the coastline. Every S-Unit
	has three components: an area of sea (the seaward component), a length
	ot coastline (the coastline component), and an area of land (the landward
<u></u>	component).
Significance	An impact is significant if, in isolation or in combination with other
	impacts, it should be taken into account in the decision-making process.
	The degree of significance depends on the combination of impact
<u></u>	magnitude and probability of impact occurrence.
Site selection	A study that was undertaken to determine the preferred site for the
process	onsnore LING Facilities. This included looking at a number of locations
	throughout Cabo Deigado Province from an environmental, social and
Class estates	engineering perspective.
Slug catcher	Liquids and gas do not form a homogeneous mixture as they flow
	through the pipeline. Liquids tend to pool in low areas of the pipeline
	this accurate processing is built on baking the liquid and the liquid in
	uns occurs, pressure is built up bening the liquid and the liquid is
	nuched through the ningline. This phonomenant is referred to a film
	pushed through the pipeline. This phenomenon is referred to as 'slug
	pushed through the pipeline. This phenomenon is referred to as 'slug flow'. A slug catcher is a device that receives the 'slugs' of liquid that accumulate over time and are pushed through the pipeline.
	pushed through the pipeline. This phenomenon is referred to as 'slug flow'. A slug catcher is a device that receives the 'slugs' of liquid that accumulate over time and are pushed through the pipeline intermittently.

Sound exposure	The total noise energy delivered over a measured duration averaged
level	over one second.
Specialist studies	The technical studies undertaken by specialists for the purpose of the
	EIA (eg marine ecology, socio-economics, etc).
Stakeholder	The process of engagement between stakeholders (the proponent,
engagement	authorities and I&APs) during the planning, assessment, implementation
	and/or management of proposals or activities.
Stakeholders	See interested and affected parties (I&APs) above.
Study Area	The entire area under baseline investigation, where impacts may occur as
	a result of the Project. This may differ between specialist studies and may
	include areas outside the Afungi Project Site. For example, the Study
	Area for the landscape and seascape visual study includes a radius of
	30km from the centre of the Afungi Project Site, as it is considered that
	the LNG Facility may be visible at such a distance. The Study Area
	comprises the ADI and AII.
Subtidal zone	The area below low tide that is almost always submerged.
Supratidal zone	The area above the high tide water line that extends upland. The area is
	seldom covered by water unless during flooding or storms, but can
	receive water from wave splash.
Surficial	The layer at the top of the sediment body, usually millimetres to
	centimetres thick.
Survey Area	The area over which physical surveys take place as part of the collection
	of primary baseline data. As in the case of Study Area, this is defined for
	each relevant resource/receptor and will differ for each specialist study,
	but these are typically confined to the Afungi Project Site or ADI.
The Project	The proposed LNG Project.
Thermocline	A layer of rapid temperature change in the water column.
Total suspended	A measure of water quality that investigates the level of suspended
solids (TSS)	solids in mg/l (turbidity).
Tungi Bay	Old name for Palma Bay.
Turbidity	The measure of the degree to which water losses transparency, due to the
<i>,</i>	presence of suspended particles.
Unexploded	Unexploded ordnances are explosive weapons (bombs, bullets, shells,
ordinance (UXO)	grenades, landmines, naval mines, etc.) that did not explode when they
ordinance (070)	were employed and still pose a risk of detonation, potentially many
	decades after they were used or discarded.
Unplanned event	A reasonably foreseeable event that is not planned to occur as part of the
emplained event	Project, but which may conceivably occur as a result of Project activities
	(eg accidents), even with a low probability.
Well	The hole drilled to obtain the gas.
Well blowout	A well blowout is an uncontrolled flow of reservoir fluids into the well
, en provout	bore and sometimes to the surface.
Wellhead	The topmost point of a well and the structure built over it
, , ciii icuu	The tophilot point of a wen and the structure built over it.